arXiv:0809.2994v3 [math.AG] 4 Dec 2008

Derived categories of small toric Calabi-Yau 3-folds and curve counting invariants

Kentaro Nagao

December 4, 2008

Abstract

We first construct a derived equivalence between a small crepant resolution of an affine toric Calabi-Yau 3-fold and a certain quiver with a superpotential. Under this derived equivalence we establish a wall-crossing formula for the generating function of the counting invariants of perverse coherent sheaves. As an application we provide some equations on Donaldson-Thomas, Pandeharipande-Thomas and Szendroi's invariants. Finally, we show that moduli spaces associated with a quiver given by successive mutations are realized as the moduli spaces associated the original quiver by changing the stability conditions.

Introduction

This is a subsequent paper of [NN]. We study *variants* of *Donaldson-Thomas* (DT in short) *invariants* on *small* crepant resolutions of affine toric Calabi-Yau varieties.

The original Donaldson-Thomas invariants of a Calabi-Yau 3-fold Y are defined by virtual counting of moduli spaces of ideal sheaves $\mathcal{I}_Z$ of 1-dimensional closed subschemes $Z \subset Y$ ([Tho00], [Beh]). These are conjecturally equivalent to *Gromov-Witten invariants* after normalizing the contribution of 0-dimensional sheaves ([MNOP06]).

A variant has been introduced Pandharipande and Thomas (PT in short) as virtual counting of moduli spaces of stable *coherent systems* ([PTa]). They conjectured these invariants also coincide with DT invariants after suitable normalization and mentioned that the coincidence should be recognized as a wall-crossing phenomenon. Here, a coherent system is a pair of a coherent sheaf and a morphism to it from the structure sheaf, which is first introduced by Le Potier in his study on moduli problems ([LP93]). Note that an ideal sheaf $\mathcal{I}_Z$ is the kernel of the canonical surjections from the structure sheaf $\mathcal{O}_Y$ to the structure sheaf $\mathcal{O}_Z$. So in this sense DT invariants also count coherent systems.

On the other hand, a variety sometimes has a derived equivalence with a noncommutative algebra. A typical example is a *noncommutative crepant resolution* of a Calabi-Yau 3-fold introduced by Michel Van den Bergh ([VdB04], [VdB]). In the case of [VdB04], the Abelian category of modules of the noncommutative crepant resolution corresponds to the Abelian category of *perverse coherent sheaves* in the sense of Tom Bridgeland ([Bri02]). Recently, Balazs Szendroi proposed to study counting invariants of ideals of such noncommutative algebras

([Sze]). He call these invariants *noncommutative Donaldson-Thomas* (NCDT in short) *invariants*. He originally studied on the conifold, but his definition works in more general settings ([Youb], [MR]).

Inspired by his work, Hiraku Nakajima and the author introduced *perverse coherent systems* (pairs of a perverse coherent sheaf and a morphism to it from the structure sheaf) and study their moduli spaces and counting invariants ([NN]). This attempt seems successful since

- we can describe explicitly a space of stability parameters with a chamber structure, and
- at certain chambers, the moduli spaces in DT, PT and NCDT theory are recovered.

Moreover, in the conifold case, we established the wall-crossing formula for the generating functions of counting invariants of perverse coherent systems and provide some equations on DT, PT and NCDT invariants. The chamber structure and the wall-crossing formula formally look very similar to the counter parts for moduli spaces of perverse coherent sheaves on the blow-up of a complex surface studied earlier by Nakajima and Yoshioka [NYa, NYb].

The purpose of this paper is to show the wall-crossing formula (Theorem 2.18) for general small crepant resolutions of toric Calabi-Yau 3-folds. Here we say a crepant resolutions of affine toric Calabi-Yau 3-fold is *small* when the dimensions of the fibers are less than 2. In such cases, the lattice polygon in $\mathbb{R}^2$ corresponding to the affine toric Calabi-Yau 3-fold does not have any lattice points in its interior. Such lattice polygons are classified up to equivalence into the following two cases:

- trapezoids with heights 1, or
- the right isosceles triangle with with length 2 isosceles edges.

In this paper we study the first case. Our argument works for the second case as well.

In §1, we construct derived equivalences between small crepant resolutions of affine toric Calabi-Yau 3-folds and certain quivers with superpotentials. In §1.1, using toric geometry, we construct tilting vector bundles given by Van den Bergh ([VdB04]) explicitly. Then, we review Ishii and Ueda's construction of crepant resolutions as moduli spaces of representations of certain quivers with superpotentials ([IU]) in §1.2. In §1.3 we show the tautological vector bundles on the moduli spaces coincide with the tilting bundles given in §1.1. Using such moduli theoretic description, we calculate the endomorphism algebras of the tilting bundles in §1.4.

The argument in §2 is basically parallel to [NN]. In our case, the fiber on the origin of the affine toric variety is the type A configuration of $(-1,-1)$- or $(0,-2)$-curves. A wall in the space of stability parameters is a hypersurface which is perpendicular to a root vector of the root system of type $\hat{A}$. Stability parameters in chambers adjacent to the wall corresponding to the imaginary root realize DT theory and PT theory ([NN, §2]). Note that the story is completely parallel to that of type $\hat{A}$ quiver varieties (of rank 1), which are the moduli spaces of framed representations of type $\hat{A}$ preprojective algebras ([Nak94], [Nak98], [Nak01]). Quiver varieties associated with a stability parameter in a chamber adjacent to the imaginary wall realize Hilbert schemes of points on the minimal resolution of the Kleinian singularitie of type A, whose exceptional fiber is the type A configurations of (-2)-curves ([Nak99], [Kuz]).

Our main result is the wall-crossing formula for the generating functions of the Euler characteristics of the moduli spaces (Theorem 2.18). The contribution of a wall depends on the information of self-extensions of stable objects on the wall. Note that in the conifold case ([NN]) every wall has a single stable object on it and every stable object has a trivial self-extension. Computations of self-extensions are done in §2.6.

Note that the sets of torus fixed points on the moduli spaces in DT, PT and NCDT theory are isolated, and we can show that DT, PT and NCDT invariants coincide with the Euler characteristics of the moduli spaces. In particular, the wall-crossing formula provides a product expansion formula of the generating functions of PT invariants. The indices in this formula are nothing but the BPS state counts $n_{g,\beta}$ ([GV], [HST01], [Tod]) in the sense of Pandeharipande-Thomas ([PTa, §3]). Although an algorithm to extract Gopakumar-Vafa invariants of our toric Calabi-Yau 3-folds from the topological vertex expression is known ([IKP] [1]), the explicit formula in this paper is new as far as the author knows.

In §3, we provide alternative descriptions of the moduli spaces. Given a quiver with a superpotential $A = (Q, \omega)$, we can *mutate* it at a vertex k to provide a new quiver with a superpotential $\mu_k(A) = (\mu_k(Q), \mu_k(\omega))$. For a generic stability parameter ζ, we can associate a sequence $k_1, \dots k_r$ of vertices and the moduli space of ζ-stable A-modules is isomorphic to the moduli space of cyclic modules over the quiver with the potetial $\mu_{k_r} \circ \cdots \circ \mu_{k_1}(A)$. As an application, we show that for a stability parameter "between DT and NCDT" the set of torus fixed points on the moduli space is isolated.

As in [NN], our formula does not cover the wall corresponding to the DT-PT conjecture. We can provide the wall-crossing formula for this wall applying Joyce's formula ([Joy]) [2]. In §4, we make some observations on how the virtual counting version of the wall-crossing formula would be induced from the perspective of the recent work of Kontsevich and Soibelman ([KS]). In fact, the wall-crossing formula (and hence the invariants) coincides with the Euler characteristic version up to sign. The author learned from Tom Bridgeland that he and Balazs Szendroi reproved the Young's product formula ([Youa]) for NCDT invariants of the conifold. Their idea looks quite similar to our observation.

Acknowledgement

The author is grateful to Hiraku Nakajima for collaborating in the paper [NN] and for many valuable discussion.

He thanks Kazushi Ueda for patiently teaching his work on brane tilings, Tom Bridgeland for explaining his work with Balazs Szendroi, Yan Soibelman for helpful comments on §4.

He also thanks Yoshiyuki Kimura and Michael Wemyss for useful discussions, Akira Ishii, Yukari Ito, Osamu Iyama, Yukiko Konishi, Sergey Mozgovoy

---

[1] The author was informed on this reference by Yukiko Konishi.

[2] Yukinobu Toda informed me that it is possible to prove the (Euler characteristic version of) DT-PT correspondence conjecture for arbitrary projective Calabi-Yau 3-folds using Joyce's formula.

and Yukinobu Toda for helpful comments. The author is supported by JSPS Fellowships for Young Scientists No.19-2672.

1 Derived equivalences

1.1 tilting generators

Let $N_0 > 0$ and $N_1 \geq 0$ be integers such that $N_0 \geq N_1$ and set $N = N_0 + N_1$. We set

$$\begin{aligned} I &= \{1, \dots, N-1\}, \\ \hat{I} &= \{0, 1, \dots, N-1\}, \\ \tilde{I} &= \left\{\frac{1}{2}, \frac{3}{2}, \dots, N - \frac{1}{2}\right\}, \\ \tilde{Z} &= \left\{ n + \frac{1}{2} \,\middle|\, n \in \mathbb{Z} \right\}. \end{aligned}$$

For $l \in \mathbb{Z}$ and $j \in \tilde{\mathbb{Z}}$, let $\underline{l} \in \hat{I}$ and $\underline{j} \in \tilde{I}$ be the elements such that $l - \underline{l} \equiv j - \underline{j} \equiv 0$ modulo N.

We denote by Γ the quadrilateral (or the triangle in the cases $N_1 = 0$) in $\mathbb{R}^2 = \{(x, y)\}$ with vertices $(0, 0)$, $(0, 1)$, $(N_0, 0)$ and $(N_1, 1)$. Let $M^\vee \simeq \mathbb{Z}^3$ be the lattice with basis $\{x^\vee, y^\vee, z^\vee\}$, and we identify the plane

$$\{(x, y, 1)\} \subset M_{\mathbb{R}}^\vee := M^\vee \otimes \mathbb{R}$$

with the one where the quadrilateral Γ is. Let M be the dual lattice of $M^\vee$. We denote the cone of Γ in $M_{\mathbb{R}}^\vee$ by Δ and consider the semigroup

$$S_\Delta = \Delta^\vee \cap M := \{u \in M \mid \langle u, v \rangle \geq 0 \ (\forall v \in \Delta)\}.$$

Let $R = R_\Gamma := \mathbb{C}[S_\Delta]$ be the group algebra and $X = X_\Gamma := \mathrm{Spce}(R_\Gamma)$ the 3-dimensional affine toric Calabi-Yau variety corresponding to Δ.

Let $\{x, y, z\} \subset M$ be the dual basis. The semigroup is generated by

$$X := x, \tag{1}$$

$$Y := -x - (N_0 - N_1)y + N_0 z, \tag{2}$$

$$Z := y, \tag{3}$$

$$W := -y + z, \tag{4}$$

and they have a unique relation $X + Y = N_0 Z + N_1 W$. So we have

$$R \simeq \mathbb{C}[X, Y, Z, W]/(XY - Z^{N_0} W^{N_1}).$$

A partition σ of Γ is a pair of functions $\sigma_x \colon \tilde{I} \to \tilde{\mathbb{Z}}$ and $\sigma_y \colon \tilde{I} \to \{0, 1\}$ such that

- $\sigma(i) := (\sigma_x(i), \sigma_y(i))$ gives a permutation of the set

$$\left\{ \left(\frac{1}{2}, 0\right), \left(\frac{3}{2}, 0\right), \dots, \left(N_0 - \frac{1}{2}, 0\right), \left(\frac{1}{2}, 1\right), \left(\frac{3}{2}, 1\right), \dots, \left(N_1 - \frac{1}{2}, 1\right) \right\},$$

- if $i < j$ and $\sigma_y(i) = \sigma_y(j)$ then $\sigma_x(i) > \sigma_x(j)$.

Giving a partition σ of Γ is equivalent dividing Γ into N-tuples of triangles $\{T_i\}_{i \in \tilde{I}}$ with area $1/2$ so that T_i has $(\sigma_x(i) \pm 1/2, \sigma_y(i))$ as its vertices. Let Γ_σ be the corresponding diagram, Δ_σ be the fun and $f_\sigma \colon Y_\sigma \to X$ be the crepant resolution of X.

We denote by $D_{\varepsilon,x}$ ($\varepsilon = 0, 1$ and $0 \leq k \leq N_\varepsilon$) the divisor of Y_σ corresponding to the lattice point (x, ε) in the diagram Γ_σ. Note that any torus equivariant divisor is described as a linear combination of $D_{\varepsilon,x}$'s. For a torus equivariant divisor D let $D(\varepsilon, x)$ denote its coefficient of $D_{\varepsilon,x}$. The support function ψ_D of D is the piecewise linear function on $|\Delta_\sigma|$ such that $\psi_D((x, \varepsilon, 1)) = -D(\varepsilon, x)$ and such that ψ_D is linear on each cone of Δ_σ. We sometimes denote the restriction of ψ_D on the plane $\{z = 1\}$ by ψ_D as well.

Definition 1.1. *For $i \in \tilde{I}$ and $k \in I$ we define effective divisors $E_i^\pm$ and $F_k^\pm$ by*

$$E_i^+ = \sum_{j=\sigma_x(i)+\frac{1}{2}}^{N_{\sigma_y(i)}} D_{\sigma_y(i),j}, \qquad F_k^+ = \sum_{i=\frac{1}{2}}^{k-\frac{1}{2}} E_i^+,$$

$$E_i^- = \sum_{j=0}^{\sigma_x(i)-\frac{1}{2}} D_{\sigma_y(i),j}, \qquad F_k^- = \sum_{i=k+\frac{1}{2}}^{N-\frac{1}{2}} E_i^-.$$

Example 1.2. *Let us consider as an example the case $N_0 = 4$, $N_1 = 2$ and*

$$(\sigma(i))_{i \in \tilde{I}} = \left(\left(\frac{7}{2}, 0 \right), \left(\frac{3}{2}, 1 \right), \left(\frac{5}{2}, 0 \right), \left(\frac{3}{2}, 0 \right), \left(\frac{1}{2}, 1 \right), \left(\frac{1}{2}, 0 \right) \right).$$

We show the corresponding diagram Γ_σ in Figure 1. The divisors are given as follows:

$$E_{\frac{1}{2}}^+ := \begin{bmatrix} 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad F_1^+ := \begin{bmatrix} 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$E_{\frac{3}{2}}^+ := \begin{bmatrix} 0 & 0 & 1 & & \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad F_2^+ := \begin{bmatrix} 0 & 0 & 1 & & \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$E_{\frac{5}{2}}^+ := \begin{bmatrix} 0 & 0 & 0 & & \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}, \quad F_3^+ := \begin{bmatrix} 0 & 0 & 1 & & \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix},$$

$$E_{\frac{7}{2}}^+ := \begin{bmatrix} 0 & 0 & 0 & & \\ 0 & 0 & 1 & 1 & 1 \end{bmatrix}, \quad F_4^+ := \begin{bmatrix} 0 & 0 & 1 & & \\ 0 & 0 & 1 & 2 & 3 \end{bmatrix},$$

$$E_{\frac{9}{2}}^+ := \begin{bmatrix} 0 & 1 & 1 & & \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad F_5^+ := \begin{bmatrix} 0 & 1 & 2 & & \\ 0 & 0 & 1 & 2 & 3 \end{bmatrix},$$

$$E_{\frac{11}{2}}^+ := \begin{bmatrix} 0 & 0 & 0 & & \\ 0 & 1 & 1 & 1 & 1 \end{bmatrix}, \quad F_6^+ := \begin{bmatrix} 0 & 1 & 2 & & \\ 0 & 1 & 2 & 3 & 4 \end{bmatrix}.$$

Here the (ε, x)-th matrix element represent the coefficient of the divisor $D_{\varepsilon,x}$.

Lemma 1.3. *(1) $\mathcal{O}_{Y_\sigma}(E_i^+ + E_i^-) \simeq \mathcal{O}_{Y_\sigma}$,*



Figure 1: Γ_σ

(2) $\mathcal{O}_{Y_\sigma}(F_k^+) \simeq \mathcal{O}_{Y_\sigma}(F_k^-)$.

Proof. We have

$$\psi_{E_i^+ + E_i^-} = \begin{cases} y - z & (\sigma_x(i) = 0), \\ -y & (\sigma_x(i) = 1), \end{cases}$$

so the equation (1) follows. Now, for the equation (2) it is enough to show that the deviser

$$F_N^+ := \sum_{i=\frac{1}{2}}^{N-\frac{1}{2}} E_i^+$$

gives the trivial bundle. In fact, we have $\psi_{F_N^+} = -x$. ☐

We denote the line bundle $\mathcal{O}_{Y_\sigma}(F_k^+) \simeq \mathcal{O}_{Y_\sigma}(F_k^-)$ on Y_σ by L_k. We set

$$F^\pm = \sum_{k=1}^{N-1} F_k^\pm, \quad L = \bigotimes_{k=1}^{N-1} L_k.$$

Example 1.4. *In the case Example 1.2,*

$$F^+ = \begin{bmatrix} 0 & 1 & 5 & & \\ 0 & 0 & 2 & 5 & 10 \end{bmatrix}.$$

Lemma 1.5. *For $i \in \tilde{I}$ we have*

$$\begin{aligned} &F^+\left(\sigma_y(i), \sigma_x(i) + \frac{1}{2}\right) - F^+\left(\sigma_y(i), \sigma_x(i) - \frac{1}{2}\right) \\ &= F^+\left(\sigma_y(i+1), \sigma_x(i+1) + \frac{1}{2}\right) - F^+\left(\sigma_y(i+1), \sigma_x(i+1) - \frac{1}{2}\right) + 1. \end{aligned}$$

Proof. First, note that

$$E_j^+\left(\sigma_y(i), \sigma_x(i) + \frac{1}{2}\right) - E_j^+\left(\sigma_y(i), \sigma_x(i) + \frac{1}{2}\right) = \delta_{i,j}.$$

So we have

$$F_k^+\left(\sigma_y(i), \sigma_x(i) + \frac{1}{2}\right) - F_k^+\left(\sigma_y(i), \sigma_x(i) + \frac{1}{2}\right) = \begin{cases} 0 & (k < i), \\ 1 & (k > i), \end{cases}$$

and so

$$F^+\left(\sigma_y(i), \sigma_x(i) + \frac{1}{2}\right) - F^+\left(\sigma_y(i), \sigma_x(i) - \frac{1}{2}\right) = N - i - \frac{1}{2}.$$

Thus the claim follows. ☐

Proposition 1.6. *The line bundles L is generated by its global sections.*

Proof. It is enough to prove that the support function ψ_{F^+} is upper convex ([Ful93, §3.4]). It is enough to prove that ψ_{F^+} is upper convex on $T_{k-\frac{1}{2}} \cap T_{k+\frac{1}{2}}$ for any $k \in I$. We denote the edge which is the intersection of $T_{k-\frac{1}{2}}$ and $T_{k+\frac{1}{2}}$ by l_k. The configurations of l_k, $T_{k-\frac{1}{2}}$ and $T_{k+\frac{1}{2}}$ are classified into the following two cases:

(1) The union of $T_{k-\frac{1}{2}}$ and $T_{k+\frac{1}{2}}$ is a parallelogram and l_k is its diagonal. In this case, the point $(\sigma_x(k+\frac{1}{2})+\frac{1}{2}, \sigma_y(k+\frac{1}{2}))$ is the intersection of l_k and l_{k-1}, the point $(\sigma_x(k-\frac{1}{2})+\frac{1}{2}, \sigma_y(k-\frac{1}{2}))$ is the other end of l_{k-1}.

(2) The union of $T_{k-\frac{1}{2}}$ and $T_{k+\frac{1}{2}}$ is a triangle and l_k is its median line. In this case, the point $(\sigma_x(k+\frac{1}{2})+\frac{1}{2}, \sigma_y(k+\frac{1}{2}))$ is the middle point and $\sigma_x(k-\frac{1}{2}) = \sigma_x(k+\frac{1}{2})+1$, $\sigma_y(k-\frac{1}{2}) = \sigma_y(k+\frac{1}{2})$.

In both cases it follows from Lemma 1.5 that ψ_{F^+} is upper convex on $T_{k-\frac{1}{2}} \cap T_{k+\frac{1}{2}}$. □

Given a divisor D the space of global sections of the line bundle $\mathcal{O}_{Y_\sigma}(D)$ is described as follows:

$$H^0(Y_\sigma, \mathcal{O}_{Y_\sigma}(D)) \simeq \bigoplus_{u \in S^0_\Delta(D)} \mathbb{C} \cdot e_u,$$

where

$$S^0_\Delta(D) := \{u \in M \mid \langle u, v\rangle \geq \psi_D(v) \ (\forall v \in |\Delta|)\}.$$

For $u \in M$ we define

$$Z_D(u) := \{v \in |\Delta| \mid \langle u, v\rangle \geq \psi_D(v)\}.$$

Then the cohomology of the line bundle $\mathcal{O}_{Y_\sigma}(D)$ is given as follows ([Ful93, §3.5]):

$$H^k(Y_\sigma, \mathcal{O}_{Y_\sigma}(D)) \simeq \bigoplus_{u \in M} H^k(|\Delta|, |\Delta| \backslash Z_D(u); \mathbb{C}).$$

We have the exact sequence of relative cohomologies

$$\begin{array}{lllllll} 0 & \rightarrow & H^0(|\Delta|, |\Delta|\backslash Z_D(u)) & \rightarrow & H^0(|\Delta|) & \xrightarrow{i_*} & H^0(|\Delta|\backslash Z_D(u)) \\ & \rightarrow & H^1(|\Delta|, |\Delta|\backslash Z_D(u)) & \rightarrow & H^1(|\Delta|) & \longrightarrow & \cdots \end{array}$$

Note that $H^0(|\Delta|) = \mathbb{C}$, $H^1(|\Delta|) = 0$ and if $|\Delta|\backslash Z_D(u)$ is not empty then i_* does not vanish. We define

$$Z^\circ_D(u) := \{v \in |\Delta| \cap \{z = 1\} \mid \langle u, v\rangle < \psi_D(v)\},$$

then $|\Delta|\backslash Z_D(u)$ is homeomorphic to $Z^\circ_D(u) \times \mathbb{R}$.

Now, in our situation it follows from the convexity of ψ_{-F} that the number of connected components of $Z^\circ_{-F}(u)$ is at most 2. Let us denote

$$S^1_\Delta(-F) := \{u \in M \mid Z^\circ_{-F}(u) \text{ has two connected components}\}.$$

Then we have

$$H^1(Y_\sigma, \mathcal{O}_{Y_\sigma}(-F)) \simeq \bigoplus_{u \in S^1_\Delta(-F)} \mathbb{C} \cdot f_u \tag{5}$$

and the R-module structure is given by

$$e_{u'} \cdot f_u = \begin{cases} f_{u+u'} & (u + u' \in S^1_\Delta(-F)), \\ 0 & (u + u' \notin S^1_\Delta(-F)) \end{cases}$$

for $u \in S^1_\Delta(-F)$ and $u' \in S_\Delta$.

For $i \in \tilde{I}$, let $t^i_F \in M$ the element such that $\langle t^i_F, * \rangle \equiv \psi_{-F}$ on the triangle T_i. Note that $t^i_F \in S^1_\Delta(-F)$ for $i \in \left\{\frac{3}{2}, \dots, N - \frac{3}{2}\right\}$.

Proposition 1.7. *The set* $\left\{ f_{t^i_F} \,\middle|\, i \in \left\{\frac{3}{2}, \dots, N - \frac{3}{2}\right\} \right\}$ *is a set of generators of* $H^1(Y_\sigma, \mathcal{O}_{Y_\sigma}(-F))$ *as an R-module.*

Proof. It is enough to check that for any $u \in S^1_\Delta(-F)$ there exist $i \in \left\{\frac{3}{2}, \dots, N - \frac{3}{2}\right\}$ and $u' \in S_\Delta$ and such that $u = t^i_F + u'$. For $\varepsilon \in \{0, 1\}$ we put

$$m_\varepsilon := \max_{0 \le j \le N_\varepsilon} \left(\langle u, (j, \varepsilon, 1) \rangle - F(\varepsilon, j) \right) \ge 0.$$

Let $u' \in S_\Delta$ be the element such that $\langle u', (x, \varepsilon, 1) \rangle = m_\varepsilon$ for any x and ε. Note that $Z^\circ_{-F}(u)$ has two connected components, $Z^\circ_{-F}(u) \subset Z^\circ_{-F}(u - u')$ and there exist x_ε's such that $\langle u - u', (x_\varepsilon, \varepsilon, 1) \rangle - F(\varepsilon, j) = 0$. Thus $Z^\circ_{-F}(u - u')$ also has two connected components, that is, $u - u' \in S^1_\Delta(-F)$. The function $\psi_{-F} - \langle u - u', * \rangle$ is upper convex and does not take negative values on Γ. So if for some x_ε's we have $F(\varepsilon, j) - \langle u - u', (x_\varepsilon, \varepsilon, 1) \rangle = 0$, then $(x_0, 0)$ and $(x_1, 1)$ should be the end points of some edge in Γ_σ. Since $u - u' \in S^1_\Delta(-F)$, (y_0, y_1) can not be neither $(0, 0)$ nor (N_0, N_1). So $(0, y_0)$ and $(1, y_1)$ is the end points of an edge l_k for some $k \in I$. By Lemma 1.5, ψ_{-F} coincides with $\langle u - u', * \rangle$ on either T_k or T_{k+1} since otherwise $\psi_{-F} - \langle u - u', * \rangle$ takes negative values on Γ. Thus claim follows. □

For a divisor D and a effective divisor E, let $1_{D,E}$ be the canonical inclusion $\mathcal{O}_{Y_\sigma}(D) \hookrightarrow \mathcal{O}_{Y_\sigma}(D + E)$.

Let us denote effective divisors

$$G^+_i = \sum_{k=1}^{i - \frac{1}{2}} F^+_k, \quad G^-_i = \sum_{k = i + \frac{1}{2}}^{N-1} F^-_k.$$

Note that $G^+_i + G^-_i$ is linearly equivalent to F^+ by lemma 1.3 and

$$\left(\psi_{G^+_i} + \psi_{G^-_i} \right) \Big|_{T_i} \equiv 0.$$

Hence we have

$$\psi_{-F} - \langle t^i_F, * \rangle \equiv \psi_{G^+_i} + \psi_{G^-_i}.$$

Let us consider the following sequence:

$$0 \to \mathcal{O}_{Y_\sigma} \longrightarrow \mathcal{O}_{Y_\sigma}(G^+_i) \oplus \mathcal{O}_{Y_\sigma}(G^-_i) \longrightarrow \mathcal{O}_{Y_\sigma}(G^+_i + G^-_i) \to 0.$$

Here the first map is given by $1_{0, G^+_i} \oplus (-1_{0, G^-_i})$ and the second map is given by $\left(1_{G^+_i, G^-_i} \right) + \left(1_{G^-_i, G^+_i} \right)$.

Proposition 1.8. *The above sequence is exact and corresponds to the element* $f_{t_F^i} \in H^1(Y_\sigma, \mathcal{O}_{Y_\sigma}(F))$.

Proof. Let $U_j \simeq \mathbb{C}^3 = \operatorname{Spec}(\mathbb{C}[x_j, y_j.z_j])$ be the affine chart corresponding to the triangle T_j. It is enough to show that the sequence is exact on each U_j. Let P_a $(a = x, y, z)$ be the vertices of T_j. Then we have

- for $j < i$, $\psi_{G_i^+}(P_a) = \psi_{G_i^+ + G_i^-}(P_a) (=: d_{j,a})$ and $\psi_{G_i^-}(P_a) = 0$,
- for $j = i$, $\psi_{G_i^+}(P_a) = \psi_{G_i^+ + G_i^-}(P_a) = \psi_{G_i^-}(P_a) = 0$,
- for $j < i$, $\psi_{G_i^-}(P_a) = \psi_{G_i^+ + G_i^-}(P_a) (=: d_{j,a})$ and $\psi_{G_i^+}(P_a) = 0$.

The sequence in the claim is restricted on U_j to the following sequence of $\mathbb{C}[x_j, y_j.z_j]$-modules:

$$0 \to (0,0,0) \xrightarrow{1 \oplus (-1)} (0,0,0) \oplus (d_{j,x}, d_{j,y}, d_{j,z}) \xrightarrow{1+1} (d_{j,x}, d_{j,y}, d_{j,z}) \to 0.$$

Here $(0,0,0)$ (resp. $(d_{j,x}, d_{j,y}, d_{j,z})$) is spaned by

$$\{x^a y^b z^c \mid a, b, c \geq 0\} \quad (\text{resp. } \{x^a y^b z^c \mid a \geq -d_{j,x}, b \geq -d_{j,y}, c \geq -d_{j,z}\})$$

as a vector space and 1 is the map which maps $x^a y^b z^c$ to $x^a y^b z^c$. We can verify this is exact. The corresponding element in $H^1(Y_\sigma, \mathcal{O}_{Y_\sigma}(F))$ can be checked by the Čech argument in [Ful93, §3.5]. □

For $k \in I$ and $i \in \left\{\frac{3}{2}, \dots, N - \frac{3}{2}\right\}$ we define divisors

$$F_k^\Delta := F_k^+ - F_k^-, \quad H_i := \sum_{k=1}^{i-\frac{1}{2}} F_k^\Delta, \quad I_k := H_{k-\frac{1}{2}} + F_k^+,$$

and the exact sequence

$$0 \to \bigoplus_{i \in \left\{\frac{3}{2}, \dots, N-\frac{3}{2}\right\}} \mathcal{O}_{Y_\sigma}(H_i) \to \bigoplus_{k \in I} \mathcal{O}_{Y_\sigma}(I_k) \to \mathcal{O}_{Y_\sigma}(F^+) \to 0.$$

The first map the sum of compositions of the maps

$$\left(1_{H_i, F^-_{i-\frac{1}{2}}} \oplus -1_{H_i, F^+_{i+\frac{1}{2}}}\right) : \mathcal{O}_{Y_\sigma}(H_i) \longrightarrow \mathcal{O}_{Y_\sigma}(I_{i-\frac{1}{2}}) \oplus \mathcal{O}_{Y_\sigma}(I_{i+\frac{1}{2}})$$

and the canonical inclusions

$$\mathcal{O}_{Y_\sigma}(I_{i-\frac{1}{2}}) \oplus \mathcal{O}_{Y_\sigma}(I_{i+\frac{1}{2}}) \hookrightarrow \bigoplus_{k \in I} \mathcal{O}_{Y_\sigma}(I_k).$$

The second map is the sum of $1_{I_k, F^+ - I_k}$'s.

Proposition 1.9. *The above sequence gives the universal extension corresponding to the set* $\{f_{t_F^i} \mid i \in \{\frac{3}{2}, \dots, N - \frac{3}{2}\}\}$ *of generators of* $H^1(Y_\sigma, \mathcal{O}_{Y_\sigma}(F))$ *as* $H^0(Y_\sigma, \mathcal{O}_{Y_\sigma})$*-module.*

Proof. For $i \in \left\{\frac{3}{2}, \dots, N - \frac{3}{2}\right\}$ we can check the sequence

$$0 \to \mathcal{O}_{Y_\sigma}(H_i) \to \left(\oplus_k \mathcal{O}_{Y_\sigma}(I_k)\right) \big/ \left(\oplus_{j \neq i} \mathcal{O}_{Y_\sigma}(H_j)\right) \to \mathcal{O}_{Y_\sigma}(F^+) \to,$$

is isomorphic to the exact sequence in Proposition 1.8. ☐

Now we have the following theorem:

Theorem 1.10. *The direct sum* $\mathcal{O}_{Y_\sigma} \oplus \bigoplus_{k \in I} L_k$ *is a projective generator of* ${}^{-1}\mathrm{Per}(Y/X)$.

Proof. This claim follows from [VdB04, Proposition 3.2.5] and Proposition 1.9. ☐

1.2 Crepant resolutions as moduli spaces

We will associate σ with a quiver with superpotential $A_\sigma = (Q_\sigma, \omega_\sigma)$. The set of vertices of the quiver Q_σ is $\hat{I}$, which is identified with $\mathbb{Z}/N\mathbb{Z}$. The set of edges of the quiver Q_σ is given by

$$H := \left(\coprod_{i \in \tilde{I}} h_i^+ \right) \sqcup \left(\coprod_{i \in \tilde{I}} h_i^- \right) \sqcup \left(\coprod_{k \in \hat{I}_r} r_k \right).$$

Here h_i^+ (resp. h_i^-) is an edge from $i - \frac{1}{2}$ to $i + \frac{1}{2}$ (resp. from $i + \frac{1}{2}$ to $i - \frac{1}{2}$), r_k is an edge from k to itself and

$$\hat{I}_r := \left\{ k \in \hat{I} \,\middle|\, \sigma_y(k - \frac{1}{2}) = \sigma_y(k + \frac{1}{2}) \right\}.$$

The relation is given as follows:

- $h_i^+ \circ r_{i-\frac{1}{2}} = r_{i+\frac{1}{2}} \circ h_i^+$ and $r_{i-\frac{1}{2}} \circ h_i^- = h_i^- \circ r_{i+\frac{1}{2}}$ for $i \in \tilde{I}$ such that $i - \frac{1}{2}$, $i + \frac{1}{2} \in \hat{I}_r$.
- $h_i^+ \circ r_{i-\frac{1}{2}} = h_{i+1}^- \circ h_{i+1}^+ \circ h_i^+$ and $r_{i-\frac{1}{2}} \circ h_i^- = h_i^- \circ h_{i+1}^- \circ h_{i+1}^+$ for $i \in \tilde{I}$ such that $i - \frac{1}{2} \in \hat{I}_r$, $i + \frac{1}{2} \notin \hat{I}_r$.
- $h_i^+ \circ h_{i-1}^+ \circ h_{i-1}^- = r_{i+\frac{1}{2}} \circ h_i^+$ and $h_{i-1}^+ \circ h_{i-1}^- \circ h_i^- = h_i^- \circ r_{i+\frac{1}{2}}$ for $i \in \tilde{I}$ such that $i - \frac{1}{2} \notin \hat{I}_r$, $i + \frac{1}{2} \in \hat{I}_r$.
- $h_i^+ \circ h_{i-1}^+ \circ h_{i-1}^- = h_{i+1}^- \circ h_{i+1}^+ \circ h_i^+$ and $h_{i-1}^+ \circ h_{i-1}^- \circ h_i^- = h_i^- \circ h_{i+1}^- \circ h_{i+1}^+$ for $i \in \tilde{I}$ such that $i - \frac{1}{2}$, $i + \frac{1}{2} \notin \hat{I}_r$.
- $h_{i-\frac{1}{2}}^+ \circ h_{i-\frac{1}{2}}^- = h_{i+\frac{1}{2}}^- \circ h_{i+\frac{1}{2}}^+$ for $k \in \hat{I}_r$.

This quiver is derived from the following bipartite graph on a 2-dimensional torus. Let S be the union of infinite number of rhombi with edge length 1 as in Figure 2 which is located so that the centers of the rhombi are on a line parallel to the x-axis in $\mathbb{R}^2$ and H be the union of infinite number of hexagons with edge length 1 as in Figure 3 which is located so that the centers of the hexagons are in a line parallel to the x-axis in $\mathbb{R}^2$. We make the sequence $\mathbb{Z} \to \{S, H\}$ which maps l to S (resp. H) if l module N is not in $\hat{I}_r$ (resp. is in $\hat{I}_r$) and cover



Figure 2: S



Figure 3: H



Figure 4: P_σ in case Example 1.2

the whole plane $\mathbb{R}^2$ by arranging S's and H's according to this sequence (see Figure 4). We regard this as a graph on the 2-dimensional torus $\mathbb{R}^2/\Lambda$, where Λ is the lattice generated by $(\sqrt{3}, 0)$ and $(N_0 - N_1, (N_0 - N_1)\sqrt{3} + N_1)$. We can colored the vertices of this graph by black or white so that each edge connect a black vertex and a white one. Let P_σ denote this bipartite graph on the torus. For each edge $h^\vee$ in P_σ, we make its dual edge h directed so that we see the black end of $h^\vee$ on our right hand side when we cross $h^\vee$ along h in the given direction. The resulting quiver coincides with Q_σ. For each vertex q of P_σ, the composition of all arrows in Q_σ corresponding to edges in P_σ with q as their ends gives a superpotential. We define ω_σ as the sum of all such superpotentials.

Put $\delta = (1, \ldots, 1) = \mathbb{Z}^{\hat{I}}$ and take a stability parameter $\theta \in \operatorname{Hom}(\mathbb{Z}^{\hat{I}}, \mathbb{R}) \simeq \mathbb{R}^{\hat{I}}$ so that $\theta(\delta) = 0$.

Definition 1.11. *An A_σ-module V is θ-(semi)stable if for any nonzero submodule $0 \neq S \subsetneq V$ we have $\theta(\underline{\dim} S)\,(\leq)\,0$.*

By [Kin94] such a stability condition coincides with a stability condition in geometric invariant theory and we can define the moduli space $\mathfrak{M}^\sigma_\theta(\delta)$ of θ-semistable A_σ-modules V such that $\underline{\dim}\, V = \delta$. A stability parameter θ is said to be generic if θ-semistability and θ-stability are equivalent to each other.

Theorem 1.12 ([IU, Theorem 6.4])**.** *For a generic stability parameter θ, the moduli space $\mathfrak{M}^\sigma_\theta(\delta)$ is a crepant resolution of X.*

Proof. We can verify easily that the assumptions in [IU, Theorem6.4] are satisfied. We can also check that the convex hull of height changes (see [IU, §2]) is equivalent to Δ (see the description of divisors before Theorem 1.14 for example). □

Let $T \subset \mathfrak{M}_\theta(\delta)$ be the open subset consisting of representations t such that $t(h_i^\pm) \neq 0$ for any $i \in$ and $t(r_k) \neq 0$ for any $k \in \hat{I}_r$. Then T is the 3-dimensional torus acting on $\mathfrak{M}_\theta(\delta)$ by edge-wise multiplications.

1.3 Description at a specific parameter

Let θ_0 be a stability parameter so that $\theta_0(\delta) = 0$ and $(\theta_0)_k < 0$ for any $k \neq 0$.

For $i \in \tilde{I}$ let $p_i \in \mathfrak{M}_{\theta_0}(\delta)$ be the representation such that

$$p_i(h_j^+) = \begin{cases} 1 & (j < i), \\ 0 & (j \geq i), \end{cases} \quad p_i(h_j^-) = \begin{cases} 1 & (j > i), \\ 0 & (j \leq i), \end{cases} \quad p_i(r_k) = 0.$$

This is fixed by the torus action. We can take a coordinate (x_i, y_i, z_i) on the neighborhood U_i of p_i in $\mathfrak{M}^\sigma_{\theta_0}(\delta)$ such that the representation $v[x_i, y_i, z_i]$ with the coordinate (x_i, y_i, z_i) is give by

$$\begin{aligned} v[x_i, y_i, z_i](h_j^+) &= 1 \quad (j < i), \\ v[x_i, y_i, z_i](h_j^-) &= 1 \quad (j > i), \\ v[x_i, y_i, z_i](h_i^+) &= x, \\ v[x_i, y_i, z_i](h_i^-) &= y, \\ v[x_i, y_i, z_i](q^-_{i-\frac{1}{2}}) = v[x_i, y_i, z_i](q^+_{i+\frac{1}{2}}) &= z, \end{aligned}$$

where

$$q_l^{\pm} = \begin{cases} r_l & (l \in \hat{I}_r), \\ h^{\mp}_{l \pm \frac{1}{2}} \circ h^{\pm}_{l \pm \frac{1}{2}} & (l \notin \hat{I}_r). \end{cases}$$

For $k \in I$ let $c_k^1 \in \mathfrak{M}^{\sigma}_{\theta_0}(\delta)$ be the representation such that

$$\begin{aligned} c_k^1(h_j^+) &= \begin{cases} 1 & (j < k), \\ 0 & (j > k), \end{cases} \\ c_k^1(h_j^-) &= \begin{cases} 1 & (j > k), \\ 0 & (j < k), \end{cases} \\ c_k^1(r_l) &= 0, \end{aligned}$$

and C_k be the closure of the orbit of c_k^1 with respect to the torus action. This is isomorphic to $\mathbb{P}^1$, contained in $U_{k-\frac{1}{2}} \cap U_{k+\frac{1}{2}}$ and

$$C_k|_{U_{k-\frac{1}{2}}} = \{y_{k-\frac{1}{2}} = z_{k-\frac{1}{2}} = 0\}, \quad C_k|_{U_{k+\frac{1}{2}}} = \{x_{k+\frac{1}{2}} = z_{k+\frac{1}{2}} = 0\}.$$

Note that the coordinate transformation is given by

$$(x_{k+\frac{1}{2}}, y_{k+\frac{1}{2}}, z_{k+\frac{1}{2}}) \longmapsto \begin{cases} (x^2_{k-\frac{1}{2}} y_{k-\frac{1}{2}}, x^{-1}_{k-\frac{1}{2}}, z_{k-\frac{1}{2}}) & (k \in \hat{I}_r) \\ (x_{k-\frac{1}{2}} z_{k-\frac{1}{2}}, x^{-1}_{k-\frac{1}{2}}, x_{k-\frac{1}{2}} y_{k-\frac{1}{2}}) & (k \notin \hat{I}_r), \end{cases}$$

Hence C_k is a $(0,-2)$-curve if $k \in \hat{I}_r$ and a $(-1,-1)$-curve if $k \notin \hat{I}_r$.

For a crepant resolution of X the configuration of $(0,-2)$-curves and $(-1,-1)$-curves determines its toric diagrams. So we have the following proposition:

Proposition 1.13. *The crepant resolution $\mathfrak{M}^{\sigma}_{\theta_0}(\delta)$ is isomorphic to Y_σ. For $i \in \tilde{I}$ the triangle T_i corresponds to the fixed point p_i and for $k \in I$ the edge l_k corresponds to the curve C_k.*

For $\varepsilon = 0$ or 1 and $0 \le x \le N_\varepsilon$ let $d^1_{\varepsilon,x} \in \mathfrak{M}^{\sigma}_{\theta_0}(\delta)$ be the representation such that

$$\begin{aligned} d^1_{\varepsilon,x}(h_j^+) &= \begin{cases} 1 & (\sigma_y(j) \neq \varepsilon \text{ or } \sigma_x(i) > x), \\ 0 & (\text{otherwise}), \end{cases} \\ d^1_{\varepsilon,x}(h_j^-) &= \begin{cases} 1 & (\sigma_y(j) \neq \varepsilon \text{ or } \sigma_x(i) < x), \\ 0 & (\text{otherwise}), \end{cases} \\ d^1_{\varepsilon,x}(r_k) &= \begin{cases} 1 & (\sigma_y(k \pm \frac{1}{2}) = \varepsilon), \\ 0 & (\text{otherwise}), \end{cases} . \end{aligned}$$

Then the closure of the orbit of $d^1_{\varepsilon,x}$ with respect to the torus action coincides with the divisor $D_{\varepsilon,x}$.

We take the tautological line bundles L_k^{σ,θ_0} $(k \in \hat{I})$ on $\mathfrak{M}^{\sigma}_{\theta_0}(\delta)$ corresponding to the vertex k of the quiver Q_σ such that $L_0^{\sigma,\theta_0} \simeq \mathcal{O}_{\mathfrak{M}^{\sigma}_{\theta_0}(\delta)}$. We have the tautological section of the line bundle $\mathrm{Hom}(L^{\sigma,\theta_0}_{i\mp\frac{1}{2}}, L^{\sigma,\theta_0}_{i\pm\frac{1}{2}})$ corresponding to the



Figure 5: Universal representations on $\mathfrak{M}^{\sigma}_{\theta_0}(\delta)$ in case Example 1.2

edge $h_i^{\pm}$. From the description above, its divisor coincides with $E_i^{\pm}$ defined in §1.1. Hence we have

$$L_k^{\sigma,\theta_0} \simeq \mathcal{O}(\sum_{i=\frac{1}{2}}^{k-\frac{1}{2}} E_i^{+}) \simeq L_k,$$

where L_k is defined just after 1.3. In summary, we have the following theorem:

Theorem 1.14. *The tautological vector bundle* $L^{\sigma,\theta_0} = \bigoplus_{k\in\hat{I}} L_k^{\sigma,\theta_0}$ *is a projective generator of* $\mathrm{Per}(Y_\sigma/X)$.

In particular, we have the following equivalence:

$$\begin{array}{ccc} D^b(\mathrm{Coh}(Y_\sigma)) & \simeq & D^b(\mathrm{mod}(\mathrm{End}_{Y_\sigma}(L^{\sigma,\theta_0}))) \\ \cup & & \cup \\ \mathrm{Per}(Y_\sigma/X) & \simeq & \mathrm{mod}(\mathrm{End}_{Y_\sigma}(L^{\sigma,\theta_0})). \end{array}$$

1.4 Computation of the endomorphism algebra

In this subsection, we denote L_k^{σ,θ_0} and L^{σ,θ_0} simply by L_k^σ and L^σ respectively. Since L^σ is the tautological bundle on the moduli space of A_σ-modules, we have the tautological map

$$\phi\colon A_\sigma \to \mathrm{End}_{Y_\sigma}(L^\sigma).$$

Proposition 1.15. *Let* $e_k \in A_\sigma$ *be the idempotent corresponding to the vertex* k. *Then the restriction of the tautological map*

$$\phi_k\colon e_kAe_k \to \mathrm{End}_{Y_\sigma}(L_k^\sigma) \simeq H^0(Y_\sigma, \mathcal{O}_{Y_\sigma}) \simeq \mathbb{C}[X,Y,Z,W]/(XY - Z^{N_1}W^{N_0}).$$

is bijective.

Proof. We have the tautological section of the line bundle $\mathcal{E}\mathrm{nd}_{Y_\sigma}(L_k^\sigma) \simeq \mathcal{O}_{Y_\sigma}$ corresponding to the path

$$X_k := h^+_{k-\frac{1}{2}} \circ \cdots \circ h^+_{\frac{1}{2}} \circ h^+_{N-\frac{1}{2}} \circ \cdots \circ h^+_{k+\frac{3}{2}} \circ h^+_{k+\frac{1}{2}}.$$

Its divisor is

$$F_N^+ = \sum_{i=\frac{1}{2}}^{N-\frac{1}{2}} E_i^+$$

and as we have seen in Lemma 1.3 we have $-\psi_{F_N^+} = x$. So $\phi_k(X_k)$ coincides with X up to scalar multiplication (see the defining equation (1) of X in §1.1).

Similarly, we put

$$Y_k := h^-_{k+\frac{1}{2}} \circ \cdots \circ h^-_{N-\frac{1}{2}} \circ h^-_{\frac{1}{2}} \circ \cdots \circ h^-_{k-\frac{3}{2}} \circ h^-_{k-\frac{1}{2}},$$

$$Z_k := \begin{cases} h^-_{k+\frac{1}{2}} \circ h^+_{k+\frac{1}{2}} & (\sigma_y(k+\frac{1}{2}) = 1), \\ h^+_{k-\frac{1}{2}} \circ h^-_{k-\frac{1}{2}} & (\sigma_y(k-\frac{1}{2}) = 1), \\ r_k & (\text{otherwise}), \end{cases}$$

$$W_k := \begin{cases} h^-_{k+\frac{1}{2}} \circ h^+_{k+\frac{1}{2}} & (\sigma_y(k+\frac{1}{2}) = 0), \\ h^+_{k-\frac{1}{2}} \circ h^-_{k-\frac{1}{2}} & (\sigma_y(k-\frac{1}{2}) = 0), \\ r_k & (\text{otherwise}). \end{cases}$$

Then $\phi_k(Y_k)$, $\phi_k(Z_k)$ and $\phi_k(W_k)$ respectively coincide with Y, Z and W up to scalar multiplications. Hence ϕ_k is surjective. To show the injectiveness, it is enough to check that X_k, Y_k, Z_k and W_k

- generate $e_k A e_k$,
- commute with each other, and
- satisfy the relation $X_k Y_k = Z_k^{N_0} W_k^{N_1}$.

These follow from the lemma below. $\square$

Lemma 1.16. *For a path P in Q_σ starting from $k \in \tilde{I}$ let P_{red} be a minimal length path which is "homotopic" to P. Namely P_{red} is the composition of h_i^+'s or h_i^-'s and is homotopic to P when we regard Q_σ as S^1 and a path in Q_σ as a map from the interval $[0,1]$ to S^1. We set*

$$\begin{aligned} r_k(P) &= (\textit{the number of } r_k \textit{ appearing in } P), \\ h_i(P) &= (\textit{the number of } h_i^+ \textit{ appearing in } P) - (\textit{the number of } h_i^+ \textit{ appearing in } P_{\mathrm{red}}) \\ &= (\textit{the number of } h_i^- \textit{ appearing in } P) - (\textit{the number of } h_i^- \textit{ appearing in } P_{\mathrm{red}}), \end{aligned}$$

and

$$\begin{aligned} z(P) &= \sum_{\substack{l \in \hat{I}_r, \\ \sigma_y(l \pm \frac{1}{2}) = 1}} r_k(P) + \sum_{\substack{i \in \tilde{I}, \\ \sigma_y(i) = 0}} h_i(P), \\ w(P) &= \sum_{\substack{l \in \hat{I}_r, \\ \sigma_y(l \pm \frac{1}{2}) = 0}} r_l(P) + \sum_{\substack{i \in \tilde{I}, \\ \sigma_y(i) = 1}} h_i(P). \end{aligned}$$

Then we have

$$P = P_{\mathrm{red}} \circ (Z_k)^{z(P)} \circ (W_k)^{w(P)}.$$

Proof. From the relations of A_σ we can verify directly that for any path P from k to k' we have

$$Z_{k'} \circ P = P \circ Z_k, \quad W_{k'} \circ P = P \circ W_k \tag{6}$$

in A_σ.

For a path P in Q_σ from k we have a expression

$$P = P' \circ Z_{k'} \circ P'' \text{ (or } P' \circ W_{k'} \circ P'')$$

for some paths P', P'' in Q_σ and $k' \in I$ unless $P = P_{\mathrm{red}}$. This case we have

$$P = P' \circ P'' \circ Z_k \text{ (or } P \circ P'' \circ W_k)$$

in A_σ by the equation (6). Then apply the same procedure for $P' \circ P''$ unless it is reduced. By the induction with respect to the lengths of paths, we can verify the claim. $\square$

For $k \neq k' \in I$ let $X_{k,k'}$ (resp. $Y_{k,k'}$) be the minimal length path from k to k' which is a composition of h_i^+'s (resp. h_i^-'s). Then we have the following basis of $e_{k'} A_\sigma e_k$:

$$\{(X_{k'})^n \circ X_{k,k'} \circ (Z_k)^m \circ (W_k)^l\}_{n,m,l \geq 0} \sqcup \{(Y_{k'})^n \circ Y_{k,k'} \circ (Z_k)^m \circ (W_k)^l\}_{n,m,l \geq 0}.$$

We have

$$\begin{aligned} X_{k',k} \circ (X_{k'})^n \circ X_{k,k'} \circ (Z_k)^m \circ (W_k)^l &= (X_k)^{n+1} \circ (Z_k)^m \circ (W_k)^l, \\ X_{k,k'} \circ (Y_{k'})^n \circ Y_{k,k'} \circ (Z_k)^m \circ (W_k)^l &= (Y_k) \circ (Z_k)^{m+m'} \circ (W_k)^{l+l'}, \end{aligned}$$

where m' and l' is nonnegative integer such that $X_{k,k'} \circ Y_{k,k'} = (Z_k)^{m'} \circ (W_k)^{l'}$. In particular, we have

Lemma 1.17. *The map*

$$X_{k,k'} \circ - \colon e_{k'} A_\sigma e_k \to e_k A_\sigma e_k$$

is injective.

Proposition 1.18. *For $k \neq k' \in I$ the restriction of the tautological map*

$$\phi_{k,k'} \colon e_{k'} A e_k \to \mathrm{Hom}_{Y_\sigma}(L^\sigma_k, L^\sigma_{k'})$$

is bijective.

Proof. The injectiveness follows from Lemma 1.17 and the injectiveness of ϕ_k. For the surjectiveness, it is enough to check that $\mathrm{Hom}_{Y_\sigma}(L_k, L_{k'})$ is generated by the image of $X_{k,k'}$ and $Y_{k,k'}$ as an R-module. Let $F^+_{k.k'}$ (resp. $F^-_{k.k'}$) be the sum of E^+_i's (resp. E^-_i's) corresponding to $X_{k,k'}$ (resp. $Y_{k,k'}$). Note that $\psi_{F^+_{k.k'}-F^-_{k.k'}}$ is the unique element in M such that $\psi_{F^+_{k.k'}-F^-_{k.k'}}(N_\varepsilon, \varepsilon, 1) = \psi_{F^+_{k.k'}}(N_\varepsilon, \varepsilon, 1)$ and such that $\psi_{F^+_{k.k'}-F^-_{k.k'}}(1,0,0) = -1$.

Let $u \in S^0_\Delta(F^+_{k.k'})$ be an element such that $\langle u, v\rangle < 0$ for some $v \in Q$. It is enough to check there exists $u' \in S_\Delta$ such that $u = \psi_{F^+_{k.k'}-F^-_{k.k'}} + u'$. Since $\langle u, (0,0,1)\rangle, \langle u, (0,1,1)\rangle \geq 0$ and $\langle u, v\rangle < 0$ for some $v \in Q$, we have $\langle u, (1,0,0)\rangle < 0$. Let $u' \in S_\Delta$ be the element such that

$$\begin{aligned} \langle u', (1,0,0)\rangle &= \langle u, (1,0,0)\rangle + 1 \\ \langle u', (N_\varepsilon, \varepsilon, 1)\rangle &= \langle u, (N_\varepsilon, \varepsilon, 1)\rangle + F^+_{k,k'}(\varepsilon, N_\varepsilon). \end{aligned}$$

It follows from the characterization of $\psi_{F^+_{k.k'}-F^-_{k.k'}}$ above that $u-u' = \psi_{F^+_{k.k'}-F^-_{k.k'}}$. □

In summary, we have the following theorem:

Theorem 1.19. *The tautological homomorphism*

$$A_\sigma \to \mathrm{End}_{Y_\sigma}(L^\sigma).$$

is isomorphism.

2 Counting invariants

From now on, we denote Q_σ and A_σ simply by Q and A. Let Q_0 and Q_1 denote the sets of vertices and edges of the quiver Q respectively, and A-fmod denote the category of finite dimensional A-modules.

2.1 Koszul resolution

We set a grading on the path algebra $\mathbb{C}Q$ such that

$$\deg(e_k) = 0, \quad \deg(h_i^{\pm}) = 1, \quad \deg(r_k) = 2.$$

The superpotential ω is homogeneous of degree 4 with respect to this grading. So the quiver with superpotential $A = (Q, \omega)$ is graded 3-dimensional Calabi-Yau algebra.

We denote the subalgebra $\mathbb{C}Q_0 = \oplus_{k \in Q_0} \mathbb{C}e_k$ of A by S. For an S-module T we define an A-bimodule F_T by

$$F_T = A \otimes_S T \otimes_S A.$$

For $k, k' \in I$ let $T_{k,k'}$ denote the 1-dimensional S-module given by

$$e_l \cdot 1 = \delta_{k,l}, \quad 1 \cdot e_l = \delta_{k',l},$$

and we set

$$F_k := F_{T_{k,k}}, \quad F_{k,k'} := F_{T_{k,k'}}.$$

Note that an element of $F_{k,k'}$ is described as a linear combination of $\{p \otimes 1 \otimes q\}_{p \in Ae_k,\ q \in e_{k'}A}$.

For a quiver with superpotential A, the *Koszul complex* of A is the following complex of A-bimodules:

$$0 \to \bigoplus_{k \in Q_0} F_k \xrightarrow{d_3} \bigoplus_{a \in Q_1} F_{\mathrm{out}(a),\mathrm{in}(a)} \xrightarrow{d_2} \bigoplus_{b \in Q_1} F_{\mathrm{in}(b),\mathrm{out}(b)} \xrightarrow{d_1} \bigoplus_{k \in Q_0} F_k \xrightarrow{m} A \to 0.$$

Here the maps m, d_1, d_3 are given by

$$m(p \otimes 1 \otimes q) = pq \quad (p \in Ae_k,\ q \in e_kA),$$
$$d_1(p \otimes 1 \otimes q) = (pb \otimes 1 \otimes q) - (p \otimes 1 \otimes bq) \quad (p \in Ae_{\mathrm{in}(b)},\ q \in e_{\mathrm{out}(b)}A),$$
$$d_3(p \otimes 1 \otimes q) = \left(\bigoplus_{a\colon \mathrm{in}(a)=k} pa \otimes 1 \otimes q \right) - \left(\bigoplus_{a\colon \mathrm{out}(a)=k} p \otimes 1 \otimes aq \right) \quad (p \in Ae_k,\ q \in e_kA).$$

The map d_2 is defined as follows: Let c be a cycle in the quiver Q. We define the map $\partial_{c;a,b}\colon F_{\mathrm{out}(a),\mathrm{in}(a)} \to F_{\mathrm{in}(b),\mathrm{out}(b)}$ by

$$\partial_{c;a,b}(p \otimes 1 \otimes q) = \sum_{\substack{r \in e_{\mathrm{in}(a)}Ae_{\mathrm{out}(b)}, \\ s \in e_{\mathrm{in}(b)}Ae_{\mathrm{out}(a)}, \\ arbs = c}} ps \otimes 1 \otimes rq.$$

Then $d_2 = \partial_\omega$ is defined as the linear combination of ∂_c's.

Since A is graded 3-dimensional Calabi-Yau algebra, the Koszul complex is exact ([Boc08, Theorem 4.3]).

2.2 new quiver

We make a new quiver $\tilde{Q}$ by adding one vertex ∞ and one arrow from the vertex ∞ to the vertex 0 to the original quiver Q. The original superpotential

ω gives the superpotential on the new quiver $\tilde{Q}$ as well. We set $\tilde{A} := (\tilde{Q}, \omega)$ and denote the category of finite dimensional $\tilde{A}$-modules by $\tilde{A}$-fmod. Giving a finite dimensional $\tilde{A}$-module $\tilde{V}$ is equivalent to giving a pair (V, W, i) of a finite dimensional A-module V, a finite dimensional vector space W at the vertex ∞ and a linear map $i \colon W \to V_0$. Let $\iota \colon \tilde{A}\text{-fmod} \to A\text{-fmod}$ be the forgetting functor mapping (V, W, i) to V.

We also consider the following Koszul type complex $\tilde{A}$-bimodules as well:

$$0 \to \bigoplus_{k \in Q_0} \tilde{F}_k \xrightarrow{\tilde{d}_3} \bigoplus_{a \in Q_1} \tilde{F}_{\mathrm{out}(a),\mathrm{in}(a)} \xrightarrow{\tilde{d}_2} \bigoplus_{b \in \tilde{Q}_1} \tilde{F}_{\mathrm{in}(b),\mathrm{out}(b)} \xrightarrow{\tilde{d}_1} \bigoplus_{k \in \tilde{Q}_0} \tilde{F}_k \xrightarrow{\tilde{m}} \tilde{A} \to 0,$$

where $\tilde{F}_i$, $\tilde{F}_{i,i'}$, $\tilde{d}_\bullet$ and $\tilde{m}$ are defined in the same way. This is also exact. The exactness at the last three terms is equivalent to the definition of generators and relations of the algebra $\tilde{A}$. The exactness at the first two terms is derived from that of the exactness of the Koszul complex of A.

Proposition 2.1. *For E, $F \in \tilde{A}$-fmod we have*

$$\begin{aligned} &\hom_{\tilde{A}}(E,F) - \mathrm{ext}^1_{\tilde{A}}(E,F) + \mathrm{ext}^1_{\tilde{A}}(F,E) - \hom_{\tilde{A}}(F,E) \\ &= \dim E_\infty \cdot \dim F_0 - \dim E_0 \cdot \dim F_\infty. \end{aligned}$$

Proof. Note that we can compute $\mathrm{Ext}^\bullet_{\tilde{A}}(E,F)$ (resp. $\mathrm{Ext}^\bullet_A(\iota(E), \iota(F))$) using the Koszul resolution. Let $d_i^{\tilde{A}}(E,F)$ (resp. $d_i^A(\iota(E), \iota(F))$) denote the derivation in the complex derived from the Koszul resolution. Then we have

$$\begin{aligned} &\hom_{\tilde{A}}(E,F) - \mathrm{ext}^1_{\tilde{A}}(E,F) \\ = &\sum_{k \in \tilde{Q}_0} (\dim E_k \cdot \dim F_k) - \sum_{a \in \tilde{Q}_1} (\dim E_{\mathrm{out}(a)} \cdot \dim F_{\mathrm{in}(a)}) + \mathrm{rank}\left(d_2^{\tilde{A}}(E,F)\right). \end{aligned}$$

Note that

$$\begin{aligned} \mathrm{rank}\left(d_2^{\tilde{A}}(E,F)\right) &= \mathrm{rank}\left(d_2^A(\iota(E), \iota(F))\right) \\ &= \mathrm{rank}\left(d_2^A(\iota(F), \iota(E))\right) \\ &= \mathrm{rank}\left(d_2^{\tilde{A}}(F,E)\right), \end{aligned}$$

where the second equation comes from the self-duality of the Koszul complex of A. Hence we have

$$\begin{aligned} &\hom_{\tilde{A}}(E,F) - \mathrm{ext}^1_{\tilde{A}}(E,F) + \mathrm{ext}^1_{\tilde{A}}(F,E) - \hom_{\tilde{A}}(F,E) \\ = &\sum_{h \in \tilde{Q}_1} (\dim E_{\mathrm{out}(h)} \cdot \dim F_{\mathrm{in}(h)} - \dim E_{\mathrm{in}(h)} \cdot \dim F_{\mathrm{out}(h)}) \\ = &\dim E_\infty \cdot \dim F_0 - \dim E_0 \cdot \dim F_\infty. \end{aligned}$$

Here the last equation follows from the fact that for any i, $j \in I$

$$\sharp(\text{arrows from } i \text{ to } j) = \sharp(\text{arrows from } j \text{ to } i).$$

$\square$

Remark 2.2. *From the Koszul resolution of A, the last equation in the proof is equivalent to the vanishing of the Euler form on A-fmod. This is equivalent to the vanishing of the Euler form on $\mathrm{Coh}_c(Y)$. The vanishing on $\mathrm{Coh}_c(Y)$ follows from Hirzebruch-Riemann-Roch theorem.*

2.3 Counting invariants

Let $\tilde{\zeta} \in \mathbb{R}^{\tilde{Q}_0}$ be a pair of real numbers. For a finite dimensional $\tilde{A}$-module $\tilde{V}$ we set

$$\theta_{\tilde{\zeta}}(\tilde{V}) = \frac{\sum_{k\in\tilde{Q}_0} \tilde{\zeta}_k \cdot \dim \tilde{V}_k}{\sum_{k\in\tilde{Q}_0} \dim \tilde{V}_k}.$$

Definition 2.3. *A finite dimensional $\tilde{A}$-module $\tilde{V}$ is said to be $\theta_{\tilde{\zeta}}$-(semi)stable if we have*

$$\theta_{\tilde{\zeta}}(\tilde{V}')\,(\leq)\,\theta_{\tilde{\zeta}}(\tilde{V})$$

for any nonzero proper $\tilde{A}$-submodule $\tilde{V}'$

Here we adapt the convention for the short-hand notation. The above means two assertions: semistable if we have '$\leq$', and stable if we have '$<$'.

Remark 2.4. *(1) These stability conditions coincide with ones in geometric invariant theory ([Kin94]).*

(2) Given a real number c let $\tilde{\zeta}' \in \mathbb{R}^{\tilde{Q}_0}$ be the pair of real numbers $(\zeta_k+c)_{k\in\tilde{Q}_0}$. Then we have

$$\theta_{\tilde{\zeta}'}(\tilde{V}) = \theta_{\tilde{\zeta}}(\tilde{V}) + c.$$

Hence $\theta_{\tilde{\zeta}'}$-(semi)stability and $\theta_{\tilde{\zeta}}$-(semi)stability are equivalent.

Theorem 2.5 ([Rud97])**.** *Let a stability parameter $\tilde{\zeta} \in \mathbb{R}^{\tilde{Q}_0}$ be fixed.*

*(1) A finite dimensional $\tilde{A}$-module $\tilde{V} \in \tilde{A}$-*fmod *has the Harder-Narasimhan filtration :*

$$\tilde{V} = \tilde{V}_0 \supset \tilde{V}_1 \supset \cdots \supset \tilde{V}_k \supset \tilde{V}_{k+1} = 0$$

such that $\tilde{V}_i/\tilde{V}_{i+1}$ is $\theta_{\tilde{\zeta}}$-semistable for $i = 0, 1, \ldots, k$ and

$$\theta_{\tilde{\zeta}}(\tilde{V}_0/\tilde{V}_1) < \theta_{\tilde{\zeta}}(\tilde{V}_1/\tilde{V}_2) < \cdots < \theta_{\tilde{\zeta}}(\tilde{V}_k/\tilde{V}_{k+1}).$$

*(2) A finite dimensional $\theta_{\tilde{\zeta}}$-stable $\tilde{A}$-module $\tilde{V} \in \tilde{A}$-*fmod *has a Jordan-Hölder filtration :*

$$\tilde{V} = \tilde{V}_0 \supset \tilde{V}_1 \supset \cdots \supset \tilde{V}_k \supset \tilde{V}_{k+1} = 0$$

such that $\tilde{V}_i/\tilde{V}_{i+1}$ is $\theta_{\tilde{\zeta}}$-stable for $i = 0, 1, \ldots, k$ and

$$\theta_{\tilde{\zeta}}(\tilde{V}_0/\tilde{V}_1) = \theta_{\tilde{\zeta}}(\tilde{V}_1/\tilde{V}_2) = \cdots = \theta_{\tilde{\zeta}}(\tilde{V}_k/\tilde{V}_{k+1}).$$

We sometimes denote an $\tilde{A}$-module $(V, \mathbb{C}, i)$ with 1-dimensional vector space at the vertex ∞ simply by (V, i).

Definition 2.6. *(1) Given $\zeta \in \mathbb{R}^{Q_0}$, we take $\tilde{\zeta} \in \mathbb{R}^{\tilde{Q}_0}$ such that $\tilde{\zeta}_k = \zeta_k$ for $k \in Q_0$. A finite dimensional A-module V said to be ζ-(semi)stable if the $\tilde{A}$-module $(V, 0, 0)$ is θ_{ζ}-(semi)stable. The definition does not depend on the choice of $\tilde{\zeta}_\infty$.*

(2) Given $(V,i) \in \tilde{A}\text{-fmod}$ *and* $\zeta \in \mathbb{R}^{Q_0}$*, we define* $\tilde{\zeta} \in \mathbb{R}^{\tilde{Q}_0}$ *by* $\tilde{\zeta}_k = \zeta_k$ *for* $k \in Q_0$ *and*

$$\tilde{\zeta}_\infty = -\zeta \cdot \underline{\dim} V.$$

We say (V,i) *is* ζ*-(semi)stable if it is* $\theta_{\tilde{\zeta}}$*-(semi)stable.*

Lemma 2.7. *An* $\tilde{A}$*-module* (V,i) *is* ζ*-(semi)stable if the following conditions are satisfied:*

(A) for any nonzero A*-submodule* $0 \neq S \subseteq V$*, we have*

$$\zeta \cdot \underline{\dim} S \, (\leq) \, 0,$$

(B) for any proper A*-submodule* $T \subsetneq V$ *such that* $\mathrm{im}(i) \subset T_0$*, we have*

$$\zeta \cdot \underline{\dim} T \, (\leq) \, \zeta \cdot \underline{\dim} V.$$

For $\zeta \in \mathbb{R}^{Q_0}$ and $\mathbf{v} = (v_k) \in (\mathbb{Z}_{\geq 0})^{Q_0}$, let $\mathfrak{M}_\zeta(\mathbf{v})$ (resp. $\mathfrak{M}^{\mathrm{s}}_\zeta(\mathbf{v})$) denote the moduli space of ζ-semistable (resp. ζ-stable) $\tilde{A}$-modules (V,i) such that $\underline{\dim} V = \mathbf{v}$. They are constructed using geometric invariant theory ([Kin94]).

A stability parameter $\zeta \in \mathbb{R}^{Q_0}$ is said to be generic if ζ-semistability and ζ-stability are equivalent to each other. Since the defining relation of A is derived from the derivation of the superpotential, the moduli space $\mathfrak{M}_\zeta(\mathbf{v})$ has a symmetric perfect obstruction theory ([Sze, Theorem 1.3.1]). By the result of [Beh] a constructible $\mathbb{Z}$-valued function ν is defined on the moduli space $\mathfrak{M}_\zeta(\mathbf{v})$. We define the counting invariants

$$D^{\mathrm{eu}}_\zeta(\mathbf{v}) := \chi(\mathfrak{M}_\zeta(\mathbf{v})), \quad D_\zeta(\mathbf{v}) := \sum_{n \in \mathbb{Z}} n \cdot \chi(\nu^{-1}(n))$$

where $\chi(-)$ denote topological Euler numbers. We encode them into the generating functions

$$\mathcal{Z}^{\mathrm{eu}}_\zeta(\mathbf{q}) := \sum_{\mathbf{v} \in (\mathbb{Z}_{\geq 0})^{Q_0}} D^{\mathrm{eu}}_\zeta(\mathbf{v}) \cdot \mathbf{q}^{\mathbf{v}}, \quad \mathcal{Z}_\zeta(\mathbf{q}) := \sum_{\mathbf{v} \in (\mathbb{Z}_{\geq 0})^{Q_0}} D_\zeta(\mathbf{v}) \cdot \mathbf{q}^{\mathbf{v}}$$

where $\mathbf{q}^{\mathbf{v}} = \prod_{k \in Q_0} q_k^{v_k}$ and q_k's are formal variables.

The 3-dimensional torus T acts on Y, and so it also acts on $\mathfrak{M}_\zeta(\mathbf{v})$. Assume that the set of T-fixed points $\mathfrak{M}_\zeta(\mathbf{v})^T$ is isolated and finite (see §2.5 and §3.4). By the argument in the proof of [Sze, Lemma 2.5.2, Corollary 2.5.3 and Theorem 2.7.1], the contribution of a T-fixed point $P \in \mathfrak{M}_\zeta(\mathbf{v})^T$ is

$$(-1)^{v_0 + \sum_{a \in Q_1} v_{\mathrm{out}(a)} v_{\mathrm{in}(a)} - \sum_{k \in Q_0} v_k^2} = (-1)^{\sum_{k \in \hat{I}_r} v_k + \sum_{k \in I} v_k},$$

and so we have

$$\begin{aligned} \mathcal{Z}_\zeta(\mathbf{q}) &= \sum_{n \in \mathbb{Z}} (-1)^{\sum_{k \in \hat{I}_r} v_k + \sum_{k \in I} v_k} \left| \mathfrak{M}_\zeta(\mathbf{v})^T \right| \cdot \mathbf{q}^{\mathbf{v}} \\ &= \mathcal{Z}^{\mathrm{eu}}_\zeta(\mathbf{p}) \end{aligned} \tag{7}$$

where the new formal variables $\mathbf{p}$ are given by

$$p_k = \begin{cases} q_k & k \neq 0, k \in \hat{I}_r, \text{ or, } k = 0, k \neq \hat{I}_r, \\ -q_k & k \neq 0, k \neq \hat{I}_r, \text{ or, } k = 0, k \in \hat{I}_r. \end{cases}$$

2.4 Wall-crossing formula

The set of non-generic stability parameters is the union of the hyperplanes in $\mathbb{R}^{Q_0}$. Each hyperplane is called a wall and each connected component of the set of generic parameters is called a chamber. The moduli space $\mathfrak{M}_\zeta(\mathbf{v})$ does not change as long as ζ moves in a chamber.

Let $\zeta^\circ = (\zeta_k)_{k\in Q_0}$ be a stability parameter on a single wall and set $\zeta^\pm = (\zeta_k \pm \varepsilon)_{k\in Q_0}$ for sufficiently small $\varepsilon > 0$.

Proposition 2.8. *(1) Let $\tilde{V}' = (V', \mathbb{C}, i')$ be a ζ^+-stable $\tilde{A}$-module. Then we have an exact sequence*

$$0 \to \tilde{V} \to \tilde{V}' \to \tilde{V}'' \to 0,$$

where $\tilde{V} = (V, \mathbb{C}, i)$ is a ζ°-stable $\tilde{A}$-module, $\tilde{V}'' = (V'', 0, 0)$ and V'' is a ζ°-semistable A-module. The isomorphism class of $\tilde{V}$ and V'' are determined uniquely. In particular, assume $\tilde{V}' = (V', \mathbb{C}, i')$ is T-invariant then $\tilde{V}$ and V'' are also T-invariant.

(2) Let $\tilde{V}' = (V', \mathbb{C}, i')$ be a ζ^--stable $\tilde{A}$-module. Then we have an exact sequence

$$0 \to \tilde{V}'' \to \tilde{V}' \to \tilde{V} \to 0,$$

where $\tilde{V} = (V, \mathbb{C}, i)$ is a ζ°-stable $\tilde{A}$-module, $\tilde{V}'' = (V'', 0, 0)$ and V'' is a ζ°-semistable A-module. The isomorphism class of $\tilde{V}$ and V'' are determined uniquely. In particular, assume $\tilde{V}' = (V', \mathbb{C}, i')$ is T-invariant then $\tilde{V}$ and V'' are also T-invariant.

Proof. We take $\tilde{\zeta}^\circ \in \mathbb{R}^{\tilde{Q}_0}$ as in Definition 2.6. Let

$$\tilde{V}' = \tilde{V}_0 \supset \cdots \supset \tilde{V}_M \supset \tilde{V}_{M+1} = 0$$

be a Jordan-Hölder filtration of $\tilde{V}'$ with respect to the $\theta_{\tilde{\zeta}^\circ}$-stability. Since $\dim \tilde{V}'_\infty = 1$, there is an integer $0 \le m \le M$ such that $\dim(\tilde{V}_m/\tilde{V}_{m+1})_\infty = 1$ and $\dim(\tilde{V}_{m'}/\tilde{V}_{m'+1})_\infty = 0$ for any $m' \neq m$. Then for $m' \neq m$ we have

$$\zeta^+ \cdot \dim(\tilde{V}'_m/\tilde{V}_{m'+1})_1 = \varepsilon \cdot \sum_{k\in I} (\tilde{V}_{m'}/\tilde{V}_{m'+1})_k > 0.$$

From the ζ^+-stability of $\tilde{V}'$, we have $m = M$. Put $\tilde{V} = \tilde{V}_M$ and $\tilde{V}'' = \tilde{V}/\tilde{V}_M$, we have the required sequence. Note that this sequence gives a part of the Harder-Narasimhan filtration of $\tilde{V}'$ with respect to the $\theta_{\tilde{\zeta}^-}$-stability. Assume $\tilde{V}'$ is T-invariant, then it follows from the uniqueness of the Harder-Narasimhan filtration that $\tilde{V}$ and $\tilde{V}''$ are T-invariant.

We can verify the claim of (2) similarly. ☐

We identify $\mathbb{Z}^{Q_0}$ with the root lattice of affine Lie algebra of type A_N and denote the set of positive root vectors by Λ^+.

Proposition 2.9. *Assume C is a ζ-stable A-module for some $\zeta \in \mathbb{R}^{Q_0}$. Then $\underline{\dim}\, C \in \Lambda^+$. Moreover, given a positive real root $\alpha \in \Lambda^{\mathrm{re},+}$ and a stability parameter ζ° such that ζ° is on the wall W_α but not on any other wall $W_{\alpha'}$ ($\alpha' \neq \alpha$), then we have the unique ζ°-stable T-invariant A-module C such that $\underline{\dim}\, C = \alpha$.*

Proof. Note that we have the natural homomorphism

$$R \to \bigoplus_{k \in \hat{I}} \mathrm{End}_Y(L_k) \to \mathrm{End}_Y(\oplus L_k) \simeq A,$$

where the first one is the diagonal embedding. The image of this map is central subalgebra of A. Any A-module has the R-module structure given by this homomorphism. Any finite dimensional A-module is supported on finite number of points on $\mathrm{Spec}(R) = X$ and so any finite dimensional A-module C which is ζ-stable for some ζ is supported on a maximal ideal $\mathcal{I} \subset R$. Any nonzero element of $\mathcal{I}$ induces an A-module automorphism on C by the multiplication. Since C is ζ-stable, any A-module automorphism on C is either zero or isomorphic. But this can not be isomorphic, because C is supported on $\mathcal{I}$. Hence we have $\mathcal{I} \cdot C = 0$.

Suppose that $\mathcal{I}$ corresponds to a nonsingular point on X and that C is ζ-stable and $\mathcal{I} \cdot C = 0$, then we have $\underline{\dim}(C) = (1, \ldots, 1)$. The singular points on X is classified as follows:

- the unique T-invariant (X, Y, Z, W),
- $(X, Y, Z - a, W)$ $(a \neq 0)$ or
- $(X, Y, Z, W - b)$ $(b \neq 0)$.

An A-module C such that $(X, Y, Z, W) \cdot C = 0$ is a module over the preprojective algebra of type $\hat{A}_N$. The dimension of the moduli space of representations of the preprojective algebra of type $\hat{A}_N$ is $\mathbf{vCv}$, where $\mathbf{C}$ is the Cartan matrix of type $\hat{A}_N$ ([Nak94]). It there exists a ζ-stable A-module C with $\underline{\dim}\, C = \mathbf{v}$, then $\mathbf{vCv}$ is nonnegative, which is the definition of the root vectors. The uniqueness follows from the irreducibility of the moduli space ([CB02]). Let C be an A-module such that $(X, Y, Z - a, W) \cdot C = 0$ $(a \neq 0)$. For $k \in \hat{I}$ such that $\sigma_y(k + \frac{1}{2}) = 1$, $h^-_{k+\frac{1}{2}}$ and $h^+_{k+\frac{1}{2}}$ give isomorphisms between C_k and C_{k+1}. For $k \in \hat{I}$ such that $\sigma_y(k + \frac{1}{2}) = 0$, we have

$$h^+_{k+\frac{1}{2}} \circ h^-_{k+\frac{1}{2}} = 0, \quad h^-_{k+\frac{1}{2}} \circ h^+_{k+\frac{1}{2}} = 0.$$

Thus the category of A-modules C such that $(X, Y, Z - a, W) \cdot C = 0$ $(a \neq 0)$ is equivalent to the category of modules over the preprojective algebra of type $\hat{A}_{N_0}$ with $h \circ \bar{h} = 0$ for any edge h. Under this equivalence, a root vector of the root system of type $\hat{A}_{N_0}$ corresponds to a root vector of the root system of type $\hat{A}_N$. Hence for a ζ-stable A-module C such that $(X, Y, Z - a, W) \cdot C = 0$ its dimension vector is a root vector of the root system of type $\hat{A}_N$. Similarly, for a ζ-stable A-module C such that $(X, Y, Z, W - b) \cdot C = 0$ its dimension vector is a root vector of the root system of type $\hat{A}_N$. □

Remark 2.10. *Note that the fiber over the point corresponds to the maximal ideal* $(X, Y, Z - a, W)$ *(resp.* $(X, Y, Z, W - b)$*) is the* A_{N_0} *(resp.* A_{N_1}*) configuration of* $\mathbb{P}^1$*'s.*

Corollary 2.11. *The set of nongeneric parameters is the union of the hyperplanes* $W_\alpha := \{\zeta \in \mathbb{Z}^{Q_0} \mid \zeta \cdot \alpha = 0\}$ $(\alpha \in \Lambda^+)$*.*

Take a positive real root $\alpha \in \Lambda^{\mathrm{re},+}$ and a parameter $\zeta^\circ \in \mathbb{R}^{Q_0}$ which is on W_α but not on any other wall. Let C be the unique T-invariant ζ°-stable A-module such that $\zeta^\circ \cdot \underline{\dim}\, C$. We set $\zeta^\pm = (\zeta_k \pm \varepsilon)$ for sufficiently small ε. We fix these notations throughout this subsection.

Proposition 2.12 ([NN, Proposition 3.7])**.** *For a ζ°-stable $\tilde{A}$-module $\tilde{V} = (V, i)$ we have*

$$\operatorname{ext}^1_{\tilde{A}}(C, \tilde{V}) - \operatorname{ext}^1_{\tilde{A}}(\tilde{V}, C) = \dim C_0.$$

Proof. Since C and $\tilde{V}$ are ζ°-stable and not isomorphic each other, we have $\hom_{\tilde{A}}(C, \tilde{V}) = \hom_{\tilde{A}}(\tilde{V}, C) = 0$. So the claim follows form Proposition 2.1. □

Proposition 2.13.

$$\operatorname{ext}^1_A(C, C) = \begin{cases} 0 & \sum_{k \notin \hat{I}_r} \alpha_k \text{ is odd,} \\ 1 & \sum_{k \notin \hat{I}_r} \alpha_k \text{ is even.} \end{cases}$$

We prove this proposition in §2.6.

Let $\alpha \in \Lambda^{\mathrm{re},+}$ be a positive real root such that $\sum_{k \notin \hat{I}_r} \alpha_k$ is odd. In such cases, wall-crossing formulas are given in [NN]:

Theorem 2.14 ([NN, Theorem 3.9])**.**

$$\mathcal{Z}^{\mathrm{eu}}_{\zeta^-}(\mathbf{q}) = (1 + \mathbf{q}^\alpha)^{\alpha_0} \cdot \mathcal{Z}^{\mathrm{eu}}_{\zeta^+}(\mathbf{q}).$$

Remark 2.15. *To be precise we should modify the argument in [NN] a little, since the stable objects on the wall are not unique, while so are the T-invariant stable objects. See the argument after Proposition 2.17.*

We will study the case $\sum_{k \notin \hat{I}_r} \alpha_k$ is even. For a positive integer m, we have the unique indecomposable A-module C_m which is described as $m-1$ times successive extensions of C's.

Proposition 2.16. *(1) Let $\tilde{V}' = (V', \mathbb{C}, i')$ be a T-invariant ζ^+-stable $\tilde{A}$-module. Then we have an exact sequence*

$$0 \to \tilde{V} \to \tilde{V}' \to \oplus_{m' \geq 1} (C_{m'})^{\bigoplus n_{m'}} \to 0,$$

where $\tilde{V} = (V, \mathbb{C}, i)$ is a T-invariant ζ°-stable $\tilde{A}$-module. The integers $n_{m'}$ and isomorphism class of $\tilde{V}$ are determined uniquely and satisfy

$$\hom(\tilde{V}', C_m) = \sum_{m' \geq 1} n_{m'} \cdot \min(m', m).$$

Moreover, the composition of the maps

$$\mathbb{C}^{N_m} \hookrightarrow \operatorname{Hom}_{\tilde{A}}(C_m, \oplus_{m' \geq 1} (C_{m'})^{\oplus n_{m'}}) \longrightarrow \operatorname{Ext}^1_{\tilde{A}}(C, \tilde{V})$$

is injective. Here $N_m = \sum_{m' \geq m} n_{m'}$ and the first map is induced by inclusions $C_m \hookrightarrow C_{m'}$ $(m' \geq m)$. The second map is given by composing the inclusion $C \hookrightarrow C_m$ and $\tilde{V}' \in \operatorname{Ext}^1_{\tilde{A}}(\oplus_{m' \geq 1} (C_{m'})^{\oplus n_{m'}}, \tilde{V})$.

(2) *Let $\tilde{V}' = (V', \mathbb{C}, i')$ be a T-invariant ζ^--stable $\tilde{A}$-module. Then we have an exact sequence*

$$0 \to \oplus_{m' \geq 1} (C_{m'})^{\bigoplus n_{m'}} \to \tilde{V}' \to \tilde{V} \to 0,$$

where $\tilde{V} = (V, \mathbb{C}, i)$ is a T-invariant ζ°-stable $\tilde{A}$-module. The integers $n_{m'}$ and isomorphism class of $\tilde{V}$ are determined uniquely and satisfy

$$\hom(C_m, \tilde{V}') = \sum_{m' \geq 1} n_{m'} \cdot \min(m', m).$$

Moreover, the composition of the maps

$$\mathbb{C}^{N_m} \hookrightarrow \operatorname{Hom}_{\tilde{A}}(\oplus_{m' \geq 1}(C_{m'})^{\oplus n_{m'}}, C_m) \longrightarrow \operatorname{Ext}^1_{\tilde{A}}(\tilde{V}, C)$$

is injective. Here $N_m = \sum_{m' \geq m} n_{m'}$ and the first map is induced by surjections $C_{m'} \twoheadrightarrow C_m$ $(m' \geq m)$. The second map is given by composing the surjection $C_m \twoheadrightarrow C$ and $\tilde{V}' \in \operatorname{Ext}^1_{\tilde{A}}(\tilde{V}, \oplus_{m' \geq 1}(C_{m'})^{\oplus n_{m'}})$.

Proof. The existence of the sequences follow from Proposition 2.8, Proposition 2.9 and Proposition 2.13. Since $\hom(\tilde{V}, C_m) = 0$, we have $\hom(\tilde{V}', C_m) = \hom(\oplus_{m'} C_{m'}, \tilde{V}')$ from the long exact sequences. The middle equations follow from $\hom(C_{m'}, C_m) = \min(m', m)$. The compositions of the maps are injective since otherwise $\tilde{V}'$ has C as its direct summand. ☐

Given a sequence $(N_m)_{m \in \mathbb{Z}}$ such that $N_{m''} = N_{m''+1} = \cdots < N$ for some m'', let $Fl((N_k); N)$ be the flag variety

$$\{0 \subseteq W_1 \subseteq \cdots \subseteq W_{m''} \subseteq \mathbb{C}^N \mid \dim W_m = N_m\}.$$

We can verify the following claim as well:

Proposition 2.17. (1) *Let $\tilde{V} = (V, i)$ be a T-invariant ζ°-stable $\tilde{A}$-module. For an element*

$$(W_k) \in Fl((N_k); \operatorname{ext}^1_{\tilde{A}}(C, \tilde{V}))^T,$$

let $\tilde{V}'$ denote the $\tilde{A}$-module given by the universal extension

$$0 \to \tilde{V} \to \tilde{V}' \to \bigoplus_{m \geq 1} (C_m)^{\oplus n_m} \to 0,$$

such that the image of the composition map in Proposition 2.16 coincides with W_k. Then $\tilde{V}'$ is T-invariant and ζ^+-stable.

(2) *Let $\tilde{V} = (V, i)$ be a T-invariant ζ°-stable $\tilde{A}$-module. For an element*

$$(W_k) \in Fl((N_k); \operatorname{ext}^1_{\tilde{A}}(\tilde{V}, C))^T,$$

let $\tilde{V}'$ denote the $\tilde{A}$-module given by the universal extension

$$0 \to \bigoplus_{m \geq 1} (C_m)^{\oplus n_m} \to \tilde{V}' \to \tilde{V} \to 0$$

such that the image of the composition map in Proposition 2.16 coincides with W_k. Then $\tilde{V}'$ is T-invariant and ζ^--stable.

Hereafter we denote the set of T-fixed points on X by ${}^TX$. Let $R(T)$ be the repsentation ring of T. For a nonnegative integer N (resp. $\mathcal{N} \in R(T)$), let ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_N$ (resp. ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_{\mathcal{N}}$) denote the subscheme of ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})$ consisting of closed points $\tilde{V}$ such that $\operatorname{ext}^1(C,\tilde{V}) = N$ (resp. $\operatorname{Ext}^1(C,\tilde{V}) = \mathcal{N}$ as T-modules). Let ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m)}$ denote the subscheme of ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')$ consisting of closed points $\tilde{V}'$ such that $\hom(\tilde{V}', C_m) = \sum_{m'\geq 1} n_{m'} \cdot \min(m', m)$ We have the canonical morphism ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m)} \to {}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})$ where $\mathbf{v} = \mathbf{v}' - \sum_m mn_m \cdot \underline{\dim}\, C$ such that a closed point $\tilde{V}' \in {}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m)}$ is mapped to the closed point $\tilde{V} \in {}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})$ appeared in the exact sequence

$$0 \to \tilde{V} \to \tilde{V}' \to \oplus_{m'\geq 1}(C_{m'})^{\bigoplus n_{m'}} \to 0.$$

Let ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m),N}$ (resp. ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m),\mathcal{N}}$) denote the inverse image of ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_N$ (resp. ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_N$) with respect to the above morphism. Similarly, we define ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})^N$, ${}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})^{\mathcal{N}}$, ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')^{(n_m)}$, ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')^{(n_m),\mathcal{N}}$ and ${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')^{(n_m),N}$.

By Proposition 2.16 and Proposition 2.17, the natural map

$${}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m),\mathcal{N}} \to {}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_{\mathcal{N}}$$

is a fibration. So we have

$$\begin{aligned}
\sum_{\mathbf{v}'} \chi({}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')) \cdot \mathbf{q}^{\mathbf{v}'} &= \sum_{\mathbf{v}',(n_m),\mathcal{N}} \chi({}^T\mathfrak{M}_{\zeta^+}(\mathbf{v}')_{(n_m),\mathcal{N}}) \cdot \mathbf{q}^{\mathbf{v}'} \\
&= \sum_{\mathbf{v},(n_m),\mathcal{N}} \chi(Fl((N_m);N)) \cdot \chi({}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_{\mathcal{N}}) \cdot \mathbf{q}^{\mathbf{v}+\sum mn_m \cdot \underline{\dim}(C)} \\
&= \sum_{\mathbf{v},N} \left(\sum_{(n_m)} \chi(Fl((n_m);N)) \cdot \mathbf{q}^{mn_m \cdot \underline{\dim}(C)} \right) \chi({}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_N) \cdot \mathbf{q}^{\mathbf{v}} \\
&= \sum_{\mathbf{v},N} \left(1 - \mathbf{q}^{\underline{\dim}(C)}\right)^{-N} \chi({}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_N) \cdot \mathbf{q}^{\mathbf{v}}.
\end{aligned}$$

Similarly we have

$$\sum_{\mathbf{v}'} \chi({}^T\mathfrak{M}_{\zeta^-}(\mathbf{v}')) \cdot \mathbf{q}^{\mathbf{v}'} = \sum_{\mathbf{v},N} \left(1 - \mathbf{q}^{\underline{\dim}(C)}\right)^{-N} \chi({}^T\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})^N) \cdot \mathbf{q}^{\mathbf{v}}.$$

By Proposition 2.12 we have $\mathfrak{M}_\zeta(\mathbf{v})_N = \mathfrak{M}_\zeta(\mathbf{v})^{N+\dim C_0}$. Hence we have

$$\begin{aligned}
\sum_{\mathbf{v}'} \chi(\mathfrak{M}_{\zeta^-}(\mathbf{v}')) \cdot \mathbf{q}^{\mathbf{v}'} &= \sum_{\mathbf{v},N} \left(1 - \mathbf{q}^{\underline{\dim}(C)}\right)^{-N} \chi(\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})^N) \cdot \mathbf{q}^{\mathbf{v}} \\
&= \sum_{\mathbf{v},N} \left(1 - \mathbf{q}^{\underline{\dim}(C)}\right)^{-N} \chi(\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_{N-\dim C_0}) \cdot \mathbf{q}^{\mathbf{v}} \\
&= \sum_{\mathbf{v},N} \left(1 - \mathbf{q}^{\underline{\dim}(C)}\right)^{-N-\dim C_0} \chi(\mathfrak{M}^{\mathrm{s}}_{\zeta^\circ}(\mathbf{v})_N) \cdot \mathbf{q}^{\mathbf{v}} \\
&= \left(1 - \mathbf{q}^{\underline{\dim}(C)}\right)^{-\dim C_0} \cdot \sum_{\mathbf{v}'} \chi(\mathfrak{M}_{\zeta^+}(\mathbf{v}')) \cdot \mathbf{q}^{\mathbf{v}'}.
\end{aligned}$$

In summary, we have the following *wall-crossing formula*:

Theorem 2.18. *We have*

$$\mathcal{Z}^{\mathrm{eu}}_{\zeta^-}(\mathbf{q}) = (1-\varepsilon(\alpha)\mathbf{q}^{\alpha})^{-\varepsilon(\alpha)\alpha_0}\cdot\mathcal{Z}^{\mathrm{eu}}_{\zeta^+}(\mathbf{q}),$$

where

$$\varepsilon(\alpha) = (-1)^{\sum_{k\notin\hat{I}_r}\alpha_k}.$$

2.5 DT, PT and NCDT

Note that the set $\{[C_k]\}_{k\in I}$ form a basis of $H_2(Y;\mathbb{Z})$. We identify $H_2(Y)$ with $\mathbb{Z}^I$. For $n\in\mathbb{Z}_{\geq 0}$ and $\beta\in H_2(Y)$, let $I_n(Y,\beta)$ denote the moduli space of ideal sheaves $\mathcal{I}_Z$ of one dimensional subschemes $\mathcal{O}_Z\subset\mathcal{O}_Y$ such that $\chi(\mathcal{O}_Z)=n$ and $[Z]=\beta$. Note that the set of torus fixed points $I_n(Y,\beta)^T$ is isolated and finite ([MNOP06]). We define the *Donaldson-Thomas invariants* $I_{n,\beta}$ from $I_n(Y,\beta)$ using Behrend's function as is §2.3 ([Tho00], [Beh]), and their generating function by

$$\mathcal{Z}_{\mathrm{DT}}(Y;q,\mathbf{t}) := \sum_{n,\beta} I_{n,\beta}\cdot q^n\mathbf{t}^{\beta}$$

where $\mathbf{t}^{\beta}=\prod_{k\in I}t_k^{\beta_k}$ and t_k $(k\in I)$ is a formal variable.

Let $P_n(Y,\beta)$ denote the moduli space of stable pairs (F,s) such that $\chi(F)=n$ and $[\mathrm{supp}(F)]=\beta$. The set of torus fixed points $I_n(Y,\beta)^T$ is also isolated and finite ([PTb]). We define the *Pandharipande-Thomas invariants* $P_{n,\beta}$ and their generating function $\mathcal{Z}_{\mathrm{PT}}(Y;q,\mathbf{t})$ similarly ([PTa]).

We set $\zeta^{\circ}=(-N+1,1,\ldots,1)$, $\zeta^{\pm}=(-N+1\pm\varepsilon,1,\ldots,1)$ and $q=q_0\cdot q_1\cdot\cdots\cdot q_{N-1}$. By the result in [NN, §2] and the fact that the sets of torus fixed points are isolated and finite, we have the following theorem:

Proposition 2.19.

$$\mathcal{Z}_{\mathrm{DT}}(Y;q,q_1^{-1},\ldots,q_{N-1}^{-1}) = \mathcal{Z}_{\zeta^-}(\mathbf{q}),\quad \mathcal{Z}_{\mathrm{PT}}(Y;q,q_1^{-1},\ldots,q_{N-1}^{-1}) = \mathcal{Z}_{\zeta^+}(\mathbf{q}).$$

Let ζ_{triv} be a parameter such that $(\zeta_{\mathrm{triv}})_k>0$ for any k. Note that $\mathfrak{M}_{\zeta_{\mathrm{triv}}}(\mathbf{v})$ is empty unless $\mathbf{v}=0$ and so $\mathcal{Z}_{\zeta_{\mathrm{triv}}}(\mathbf{q})=1$. Let ζ_{cyclic} be a parameter such that $(\zeta_{\mathrm{cyclic}})_k<0$ for any k. The invariants $D_{\zeta_{\mathrm{cyclic}}}(\mathbf{v})$ are the *non-commutative Donaldson-Thomas invariants* defined in [Sze]. Note that the set of torus fixed points $\mathfrak{M}_{\zeta_{\mathrm{cyclic}}}(\mathbf{v})^T$ is isolated and finite. We denote their generating function $\mathcal{Z}_{\zeta_{\mathrm{cyclic}}}(\mathbf{q})$ by $\mathcal{Z}_{\mathrm{NCDT}}(\mathbf{q})$.

We divide the set of positive real roots into the following two parts:

$$\Lambda^{\mathrm{re},+}_{\pm} = \{\alpha\in\Lambda^{\mathrm{re},+}\mid \pm\zeta^{\circ}\cdot\alpha<0\}.$$

Applying the wall-crossing formula and comparing the equations (7) in §2.3, we obtain the following relations between generating functions:

Theorem 2.20.

$$\mathcal{Z}_{\mathrm{NCDT}}(\mathbf{q}) = \left(\prod_{\alpha\in\Delta^{\mathrm{re},+}_{-}}(1+(-1)^{\alpha_0}\mathbf{q}^{\alpha})^{\varepsilon(\alpha)\alpha_0}\right)\cdot\mathcal{Z}_{\mathrm{DT}}(Y;q,q_1^{-1},\ldots,q_{N-1}^{-1}),$$

$$\mathcal{Z}_{\mathrm{PT}}(Y;q,q_1^{-1},\ldots,q_{N-1}^{-1}) = \prod_{\alpha\in\Delta^{\mathrm{re},+}_{+}}(1+(-1)^{\alpha_0}\mathbf{q}^{\alpha})^{\varepsilon(\alpha)\alpha_0}.$$

Remark 2.21. *For the case* $N_1 = 0$*, the formula on* $\mathcal{Z}_{\mathrm{NCDT}}$ *and* $\mathcal{Z}_{\mathrm{DT}}$ *have been given in [Youb].*

We define the sets of positive real roots of the finite root system by

$$\Lambda^{\mathrm{fin},+} = \{\alpha_{[a,b]} := \alpha_a + \cdots + \alpha_b \mid 0 < a \leq b < N\},$$

then we have

$$\Lambda_+^{\mathrm{re},+} = \{\alpha + n\delta \mid \alpha \in \Lambda^{\mathrm{fin},+},\ n \geq 0\}.$$

Note that for $\alpha \in \Lambda$ we have $\varepsilon(\alpha + n\delta) = \varepsilon(\alpha)$. Let

$$M(x,q) = \prod_{n=1}^{\infty}(1 - xq^n)^{-n}$$

be the *MacMahon function*.

Corollary 2.22.

$$\mathcal{Z}_{\mathrm{PT}}(Y; q, q_1, \ldots, q_{N-1}) = \prod_{0<a\leq b<N} M(q_{[a,b]}, -q)^{\varepsilon(\alpha_{[a,b]})},$$

where $q_{[a,b]} = q_a \cdot \cdots \cdot q_b$.

The root lattice of the finite root system is identified with $H_2(Y)$ so that α_k corresponds to $[C_k]$. The corollary claims the *Gopakumar-Vafa BPS state counts* in genus g and class α *defined* in [PTa, §3.4] is given by

$$n_{g,\alpha} = \begin{cases} -\varepsilon(\alpha_{[a,b]}) & \alpha = \alpha_{[a,b]},\, g = 0, \\ 0 & \text{otherwise.} \end{cases}$$

2.6 Appendix

In this subsection, we prove Proposition 2.13. We can take integers n_0, n_1 and a basis $\{v_n\}_{n_0 \leq n \leq n_1}$ of C such that

$$C_k = \bigoplus_{\substack{n_0 \leq n \leq n_1, \\ n \equiv k \pmod N}} \mathbb{C} v_n,$$

$$\begin{aligned} \big(h_i^+(v_n), h_i^-(v_{n+1})\big) &= (v_{n+1}, 0), \text{ or } (0, v_n) \quad (i - \frac{1}{2} \equiv n), \\ h_i^-(v_{n_0}) &= 0, \quad (i - \frac{1}{2} \equiv n_0), \\ h_i^+(v_{n_1}) &= 0, \quad (i - \frac{1}{2} \equiv n_1), \\ r_k(v_n) &= 0, \quad (k \equiv n,\, k \in \hat{I}_r). \end{aligned}$$

Lemma 2.23. *Assume that we have* $d \in \mathbb{Z}$ *and* $j < j' \in \tilde{\mathbb{Z}}$ *such that* $n_0 \leq j + \frac{1}{2}, j + dN + \frac{1}{2}$ *and* $j' - \frac{1}{2}, j' + dN - \frac{1}{2} \leq n_1$*. If we have* $v_{j+\frac{1}{2}} \notin \mathrm{im}(h_{\underline{j}}^+)$*,* $v_{j+\frac{1}{2}+dN} \in \ker(h_{\underline{j}}^-)$*,* $v_{j'-\frac{1}{2}} \notin \mathrm{im}(h_{\underline{j'}}^-)$ *and* $v_{j+\frac{1}{2}+dN} \in \ker(h_{\underline{j'}}^+)$*, then* $d = 0$ *and* $\{j, j'\} = \{n_0, n_1\}$.

Proof. Note that $\bigoplus_{j<n<j'} \mathbb{C}v_n$ is a nonzero proper A-submodule of C and $\bigoplus_{j+dN<n<j'+dN} \mathbb{C}v_n$ is a nonzero proper A-quotient module of C. By the θ-stability of C we have

$$\sum_{j<n<j'} \theta_{\underline{n}} < 0, \qquad \sum_{j+dN<n<j'+dN} \theta_{\underline{n}} > 0.$$

This is a contradiction. $\square$

Using the Koszul complex, we can compute $\mathrm{Ext}^1_A(C,C)$ as the cohomology of the following complex:

$$\bigoplus_{a\in Q_1} \mathrm{Hom}(C_{\mathrm{out}(a)}, C_{\mathrm{in}(a)}) \xrightarrow{d_2} \bigoplus_{b\in Q_1} \mathrm{Hom}(C_{\mathrm{in}(b)}, C_{\mathrm{out}(b)}) \xrightarrow{d_1} \bigoplus_{k\in Q_0} \mathrm{Hom}(C_k, C_k).$$

For $d \in \mathbb{Z}$ and $j \in \tilde{\mathbb{Z}}$ such that both j and $j - \frac{1}{2} + dN$ (resp. $j + \frac{1}{2} + dN$) are in the interval $[n_0, n_1]$ we define an element

$$\alpha^{\pm}_{j,d} \in \mathrm{Hom}\left(C_{\mathrm{in}(h^{\pm}_{\underline{j}})}, C_{\mathrm{out}(h^{\pm}_{\underline{j}})}\right)$$

by

$$\alpha^{\pm}_{j,d}(v_n) = \delta_{n,j\pm\frac{1}{2}} v_{j\mp\frac{1}{2}+dN},$$

and for $l \in$ such that both l and $l + dN$ are in the interval $[n_0, n_1]$ we define an element

$$\beta_{l,d} \in \mathrm{Hom}\left(C_{\mathrm{in}(r_{\underline{l}})}, C_{\mathrm{out}(r_{\underline{l}})}\right)$$

by

$$\beta_{l,d}(v_n) = \delta_{n,l} v_{l+dN}.$$

We define a set J_0 by

$$J_0 = \{n \in \mathbb{Z} \mid n_0 \le n \le n_1,\, \underline{n} \in \hat{I}_r\} \sqcup \{j \in \tilde{\mathbb{Z}} \mid n_0 < j < n_1\}.$$

We define a set J_d for $d \in \mathbb{Z}_{>0}$ such that $dN \le n_1 - n_0$ by

$$\{n \in \mathbb{Z} \mid n_0 \le n \le n_1 - dN,\, \underline{n} \in \hat{I}_r\} \sqcup \{j \in \tilde{\mathbb{Z}} \mid n_0 < j < n_1 - dN\} \sqcup \{n_0 - 1/2\}$$

if $h^{+}_{\underline{(n_0-1/2)}}(v_{n_0-1+dN}) = 0$, and

$$\{n \in \mathbb{Z} \mid n_0 \le n \le n_1 - dN,\, \underline{n} \in \hat{I}_r\} \sqcup \{j \in \tilde{\mathbb{Z}} \mid n_0 < j < n_1 - dN\} \sqcup \{n_1 + 1/2\}$$

if $v_{n_1+1-dN} \notin \mathrm{im}\left(h^{+}_{\underline{(n_0+1/2)}}\right)$. We also define a set J_d for $d \in \mathbb{Z}_{<0}$ such that $-dN \le n_1 - n_0$ by

$$\{n \in \mathbb{Z} \mid n_0 \le n - dN \le n_1,\, \underline{n} \in \hat{I}_r\} \sqcup \{j \in \tilde{\mathbb{Z}} \mid n_0^d N < j < n_1\} \sqcup \{n_1 + 1/2\}$$

if $h^{-}_{\underline{(n_1+1/2)}}(v_{n_1+1+dN}) = 0$, and

$$\{n \in \mathbb{Z} \mid n_0 \le n \le n_1 - dN,\, \underline{n} \in \hat{I}_r\} \sqcup \{j \in \tilde{\mathbb{Z}} \mid n_0 < j < n_1 - dN\} \sqcup \{n_0 - 1/2 - dN\}$$

if $v_{n_0-1-dN}) \notin \mathrm{im}(h^{-}_{\underline{(n_0-1/2)}})$.

For $d \in \mathbb{Z}$ and $j \in \tilde{\mathbb{Z}} \cap J_d$ we take an element $\beta_{j,d}$ in the kernel of d_1 as follows:

- $\beta_{j,d} := \alpha^+_{j,d}$ if $v_{j+\frac{1}{2}} \notin \mathrm{im}(h^+_i)$ and $v_{j-\frac{1}{2}+dN} \in \ker(h^+_i)$,
- $\beta_{j,d} := \alpha^-_{j,d}$ if $v_{j-\frac{1}{2}} \notin \mathrm{im}(h^-_i)$ and $v_{j+\frac{1}{2}+dN} \in \ker(h^-_i)$,
- $\beta_{j,d} := \alpha^+_{j,d} + \alpha^-_{j,d}$ if $h^-_i(v_{j+\frac{1}{2}}) = h^+_i(v_{j-\frac{1}{2}+dN}) = 0$ or $h^+_i(v_{j-\frac{1}{2}}) = h^-_i(v_{j+\frac{1}{2}+dN}) = 0$.

Note that the set

$$\{\beta_{j,d}\}_{d\in\mathbb{Z}, j\in\tilde{\mathbb{Z}}\cap J_d} \sqcup \{\beta_{l,d}\}_{d\in\mathbb{Z}, l\in\mathbb{Z}\cap J_d}$$

forms a basis of $\ker(d_1)$.

For an element $a \in J_d$ let a^+ (resp. a^-) denote the minimal (resp. maximal) element in J_d which is larger (resp. smaller) than a if it exists. We divide J_d into two disjoint subset J'_d and J''_d so that exactly one of a or a^+ is in J'_d for any $a \in J_d$ such that a^+ exists. Furthermore we divide J'_d (resp. J''_d) into disjoint subsets satisfying the following condition: Take $a \in J'_d$ such that $b := (a^+)^+$ exists. Let H be the subset of J'_d containing a. Then $b \notin H$ if

- $a \in \tilde{\mathbb{Z}}$, $v_{a+1/2} \notin \mathrm{im}(h^+_{\underline{a}})$ and $h^-_{\underline{a}}(v_{a+1/2+dN}) = 0$, or
- $a^+ \in \tilde{\mathbb{Z}}$, $v_{a^+ +1/2} \notin \mathrm{im}(h^+_{\underline{a+}})$ and $h^-_{\underline{a+}}(v_{a^+ +1/2+dN}) = 0$, or
- $a^+ \in \tilde{\mathbb{Z}}$, $v_{a^+ -1/2} \notin \mathrm{im}(h^-_{\underline{a+}})$ and $h^+_{\underline{a+}}(v_{a^+ -1/2+dN}) = 0$, or
- $b \in \tilde{\mathbb{Z}}$, $v_{b-1/2} \notin \mathrm{im}(h^-_{\underline{b}})$ and $h^+_{\underline{b}}(v_{b-1/2+dN}) = 0$,

and otherwise $b \in H$. Note that J'_0 and J''_0 are divided into themselves.

Let H be one of the subsets of J_d ($d \neq 0$).

Lemma 2.24. *The subspace of* $\ker(d_1)$ *spanned by* $\{\beta_{a,d}\}_{a\in H}$ *is contained in* $\mathrm{im}(d_2)$.

Proof. For $a \in J_d$ we define an element $\gamma_{a,d}$ by

- $\beta_{a,d}$ if $a \in \mathbb{Z}$,
- $\bar{\alpha}^+_{a,d} \in \mathrm{Hom}\left(C_{\mathrm{out}(h^+_{\underline{a}})}, C_{\mathrm{in}(h^+_{\underline{a}})}\right)$ if $a \in \tilde{\mathbb{Z}}$ and $h^+_{\underline{a}}(v_{a-1/2}) = h^+_{\underline{a}}(v_{a+dN-1/2}) = 0$,
- $\bar{\alpha}^-_{a,d} \in \mathrm{Hom}\left(C_{\mathrm{out}(h^-_{\underline{a}})}, C_{\mathrm{in}(h^-_{\underline{a}})}\right)$ if $a \in \tilde{\mathbb{Z}}$ and $h^-_{\underline{a}}(v_{c+1/2}) = h^-_{\underline{a}}(v_{a+dN+1/2}) = 0$,

where $\bar{\alpha}^\pm_{a,d}$ is given by

$$\bar{\alpha}^\pm_{a,d}(v_n) = \delta_{n,a\mp\frac{1}{2}} v_{j+dN\pm\frac{1}{2}}.$$

For $a \in J_d$ such that a^+, $a^- \in J_d$ we have

$$d_2(\gamma_{a,d}) = \beta_{a^-,d} + \beta_{a^+,d}.$$

Let $a_{\min}$ and $a_{\max}$ be the maximal and minimal elements in H. We can verify that either $\beta_{a_{\min},d} \in \mathrm{im}(d_2)$ or $\beta_{a_{\max},d} \in \mathrm{im}(d_2)$ holds by case-by-case argument. For example, if $a_{\min} \in \mathbb{Z}$ and $h^-_{\underline{a_{\min}-}}(v_{a_{\min}^- +1/2}) = v_{a_{\min}^- -1/2}$ then we have

$$d_2(\bar{\alpha}^+_{a_{\min}^-,d}) = \beta_{a_{\min},d}.$$

□

Proposition 2.25. *If* $|J_0|$ *is even, then* J_0 *is contained in* $\mathrm{im}(d_2)$*. If* $|J_0|$ *is odd, then we have* $\dim(J_0/\mathrm{im}(d_2) \cap J_0) = 1$*.*

Proof. Let $a_{\min}$ and $a_{\max}$ be the maximal and minimal elements in J_0. For $a \in J_0$ we have

$$d_2(\gamma_{a,0}) = \begin{cases} \beta_{a_{\min}^+,d} & \text{if } a = a_{\min}, \\ \beta_{a_{\max}^-,d} & \text{if } a = a_{\max}, \\ \beta_{a^-,d} + \beta_{a^+,d} & \text{otherwise.} \end{cases}$$

Thus the claim follows. $\square$

Now we finish the proof of Proposition 2.13.

3 Mutations and stabilities

3.1 mutations

In §1.2 we associate a quiver with a potential $A := A_\tau$ with a map $\tau \colon \hat{I} \to \{\pm 1\}$, where we use 1 and −1 instead of H and S respectively. For $k \in \hat{I}$, let $\mu_k(\tau) \colon \hat{I} \to \{\pm 1\}$ be the map given by

$$\mu_k(\tau)(l) = \begin{cases} \tau(k)\tau(l) & (l = k \pm 1), \\ \tau(l) & (\text{otherwise}), \end{cases}$$

and let $\mu_k(A)$ denote the quiver with the potential $A_{\mu_k(\tau)}$.

Let P_k be the projective A-module associated with the vertex $k \in \hat{I}$ and we set $P := \bigoplus_k P_k (= A)$. We define the new A-module

$$P'_k := \mathrm{coker}(P_k \to P_{k-1} \oplus P_{k+1})$$

where $P_k \to P_{k\pm 1}$ be the maps given by conposing the arrows from $k \pm 1$ to k.

Proposition 3.1. *(1) The object* $\mu_k(P) = \bigoplus_{l \neq k} P_l \oplus P'_k$ *is a tilting generator in* $D^b(A\text{-mod})$*.*

(2) $\mathrm{End}_A(\mu_k(P)) \simeq \mu_k(A)$*.*

Proof. We provide just a skech of the proof.

For the claim (1), it is clear that $\mu_k(P)$ is a generator. Vanishing of the higher extensions from P_l $(l \neq k)$ to P'_k follows from the surjectiveness of the map

$$\mathrm{Hom}(P_l, P_{k-1}) \oplus \mathrm{Hom}(P_l, P_{k+1}) \to \mathrm{Hom}(P_l, P_k).$$

We can verify the other conditions as well.

For the claim (2), we can construct the isomorphism explicitly. Let $H_i^\pm$ and R_k denote the arrows of the mutated quiver $\mu_k(A)$, where $h_i^\pm$ and r_k are the arrows of the original quiver A. Given $k \in \hat{I}_r$, we associate the map induced by

$$(h^+_{k+\frac{1}{2}} \circ h^+_{k-\frac{1}{2}}) + (h^+_{k+\frac{1}{2}} \circ h^-_{k+\frac{1}{2}}) \colon P_{k-1} \oplus P_{k+1} \to P_{k+1}$$

with the arrow $H^+_{k+\frac{1}{2}}$, the map induced by

$$0 \oplus \mathrm{id}_{P_{k+1}} \colon P_{k+1} \to P_{k-1} \oplus P_{k+1}.$$

with the arrow $H^-_{k+\frac{1}{2}}$ and the map induced by

$$\begin{tikzcd}
P_k \arrow[r] \arrow[d, "r_k"'] & P_{k-1} \oplus P_{k+1} \arrow[d, "\begin{pmatrix} R(k-1) & 0 \\ 0 & R(k+1) \end{pmatrix}"] \\
P_k \arrow[r] & P_{k-1} \oplus P_{k+1}
\end{tikzcd}$$

with the arrow R_k. Here

$$R(k \pm 1) = \begin{cases} r_{k\pm 1} & (k \pm 1 \in \hat{I}_r), \\ h^{\mp}_{k\pm\frac{2}{3}} \circ h^{\pm}_{k\pm\frac{2}{3}} & (k \pm 1 \notin \hat{I}_r). \end{cases}$$

We can associate maps with the other arrows and verify that the construction actually provides an isomorphism. $\square$

In particular, we have the derived equivalence

$$\Psi_k \colon D^b(A\text{-mod}) \to D^b((\mu_k(A))\text{-mod}).$$

3.2 the affine Weyl group action

For $k \in \hat{I}$ we define the map $\mu_k \colon \mathbb{Z}^{\hat{I}} \to \mathbb{Z}^{\hat{I}}$ by

$$(\mu_k(\mathbf{v}))_l = \begin{cases} \mathbf{v}_{k-1} - \mathbf{v}_k + \mathbf{v}_{k+1} & l = k, \\ \mathbf{v}_l & \text{otherwise} \end{cases}$$

for $\mathbf{v} \in \mathbb{Z}^{\hat{I}}$. We also define $\mu_k \colon \mathbb{R}^{\hat{I}} \to \mathbb{R}^{\hat{I}}$ by

$$(\mu_k(\zeta))_l = \begin{cases} -\zeta_k & l = k, \\ \zeta_{k\pm 1} + \zeta_k & l = k \pm 1, \\ \zeta_l & \text{otherwise} \end{cases}$$

for $\zeta \in \mathbb{R}^{\hat{I}}$. Note that

$$\mathbf{v} \cdot \zeta = \mu_k(\mathbf{v}) \cdot \mu_k(\zeta)$$

for any $\mathbf{v}$ and ζ.

Proposition 3.2. *Let $\alpha \in \mathbb{Z}^{\hat{I}}$ be a positive root and ζ be a parameter such that $\alpha \cdot \zeta = 0$, $\zeta_k < 0$ and such that ζ is not on any other wall. Given a ζ-stable A-module C with $\underline{\dim}\, C = \alpha$ and $\mathcal{I} \cdot C = 0$ for a maximal ideal $\mathcal{I} \subset R$. Then $\Psi_k(C)$ is the unique $\mu_k(\zeta)$-stable $\mu_k(A)$-module with $\underline{\dim}(\Psi_k(C)) = \mu_k(\alpha)$ such that $\mathcal{I} \cdot \Psi_k(C) = 0$.*

Proof. Since C is ζ-stable and $\zeta_k < 0$, the map

$$C_{k-1} \oplus C_{k+1} \to C_k$$

is surjective. Hence $\Psi_k(C)$ is a $\mu_k(A)$-module. Suppose we have an exact sequence

$$0 \to V \to \Psi_k(C) \to V' \to 0$$

of nontrivial $\mu_k(A)$-modules. Let $H^*_A(-)$ denote the cohomology with respect to the natural t-structure of $D^b(A\text{-mod})$. Note that for a $\mu_k(A)$-module V, $H^*_A(\Psi_k^{-1}(V))$ is concentrated on the degree 0 and -1 and $(H_A^{-1}(\Psi_k^{-1}(V)))_l = 0$ for $l \neq k$. By the long exact sequence, we have $H_A^{-1}(\Psi_k^{-1}(V)) = 0$ and the following exact sequence

$$0 \to H_A^{-1}(\Psi_k^{-1}(V')) \to \Psi_k^{-1}(V) \to C \to H_A^0(\Psi_k^{-1}(V')) \to 0$$

of A-modules. Then we have

$$\begin{aligned}\mu_k(\zeta)\cdot \underline{\dim}\, V &= \zeta \cdot \underline{\dim}\, \Psi_k^{-1}(V)\\ &< \zeta \cdot \underline{\dim}\, H_A^{-1}(\Psi_k^{-1}(V'))\\ &\leq 0\end{aligned}$$

where the first inequality follows from the ζ-stablility of C and the second one follows from the assumption $\zeta_k < 0$. □

In the rest of this section, we fix a parameter ζ such that $\sum \zeta_l < 0$ and such that $\zeta + d\cdot \mathbf{1}$ is not on an intersetion of two walls. See Remark 3.9 for the case $\sum \zeta_l > 0$. Then we get the sequence $C_0, \dots, C_r$ of chambers such that

- $\zeta - d\cdot\mathbf{1} \in \cup \overline{C_s}$ for any $d \geq 0$,
- for any C_s, there exists some $d \geq 0$ such that $\zeta - d\cdot \mathbf{1} \in C_s$, and
- suppose $\zeta - d\cdot\mathbf{1} \in C_s$, $\zeta - d'\cdot\mathbf{1} \in C_{s'}$ and $s < s'$, then $d > d'$.

We also define the sequence $k_1, \dots, k_r$ of elements in $\hat{I}$ such that

$$\overline{C_{s-1}} \cap \overline{C_s} \subset W_{\alpha^s} \quad (\alpha^s = \mu_{k_{s-1}} \circ \cdots \circ \mu_{k_1}(\alpha_{k_s})),$$

where α_k denote the simple root vector. We denote $\mu_s := \mu_{k_s} \circ \cdots \circ \mu_{k_1}$, $\Psi_s := \Psi_{k_s} \circ \cdots \circ \Psi_{k_1}$ and $\mu_\zeta := \mu_r$, $\Psi_\zeta := \Psi_r$.

Corollary 3.3. *(1) Let C be a finite dimensional θ_ζ-stable A-module. Suppose $\underline{\dim}(C)\cdot\zeta < 0$, then $\Psi_\zeta(C) \in (\mu_\zeta(A))\text{-mod}$. Suppose $\underline{\dim}(C)\cdot\zeta > 0$, then $\Psi_\zeta(C) \in (\mu_\zeta(A))\text{-mod}[1]$.*

(2) Let C' be a finite dimensional $\theta_{\mu_\zeta(\zeta)}$-stable $\mu_\zeta(A)$-module. Take $d \in \mathbb{R}$ such that $(\mu_\zeta(\zeta) + d\cdot\mathbf{1}) \in \mu_\zeta(C^+)$ where $C^+ := \{\zeta \mid \zeta_l > 0\}$. Suppose $\underline{\dim}(C')\cdot(\mu_\zeta(\zeta) + d\cdot\mathbf{1}) > 0$, then $\Psi_\zeta^{-1}(C') \in A\text{-mod}$. Suppose $\underline{\dim}(C')\cdot(\mu_\zeta(\zeta) + d\cdot\mathbf{1}) < 0$, then $\Psi_\zeta^{-1}(C') \in A\text{-mod}[-1]$.

3.3 mutations and change of stability parameters

We set $\mathcal{P} := A$-fmod and denote by $\mathcal{P}_\zeta$ the image of the Abelian category $\mu_\zeta(A)$-fmod under the equivalence Ψ_ζ^{-1}. Let $\overline{\mathcal{P}}$ (resp. $\overline{\mathcal{P}_\zeta}$) be the category consisting of pairs (V, W, s), where $V \in \mathcal{P}$ (resp, $\in \mathcal{P}_\zeta$), W is a finite dimansional vector space and $s\colon P_0 \otimes W \to V$. An object (V, W, s) with 1-dimensional W is simply written (F, s). Note that $\overline{\mathcal{P}}$ is equivalent to $\tilde{A}$-fmod.

Definition 3.4. *For $\xi \in \mathbb{R}^{\hat{I}}$, we say an object $(V, s) \in \mathcal{P}_\zeta$ is $(\xi, \mathcal{P}_\zeta)$-(semi)stable if the following conditions are satisfied:*

(A) for any nonzero subobject $0 \neq S \subseteq V$ in $\mathcal{P}_\zeta$, we have

$$\xi \cdot \underline{\dim}\, S \,(\leq)\, 0,$$

(B) for any proper subobject $T \subsetneq V$ in $\mathcal{P}_\zeta$ which s factors through, we have

$$\xi \cdot \underline{\dim}\, T \,(\leq)\, \xi \cdot \underline{\dim}\, V.$$

From now on, the ζ-(semi)stability for objects in $\overline{\mathcal{P}} \simeq \tilde{A}$-fmod is written as the "$(\zeta, \overline{\mathcal{P}})$-(semi)stability". We set $\xi_{\text{cyclic}} := \mu_\zeta(\zeta)$. Note that $(\xi_{\text{cyclic}})_l < 0$ for any $l \in \hat{I}$. The following four claims follow from Corollary 3.3 by the same argument as in [NN, §4.3]:

Lemma 3.5. (see [NN, Lemma 4.2]) *Let (F, s) be a $(\zeta, \overline{\mathcal{P}})$-stable object, then $F \in \mathcal{P}_\zeta$.*

Proposition 3.6. (see [NN, Proposition 4.3]) *Let (F, s) be a $(\zeta, \overline{\mathcal{P}})$-stable object, then (F, s) is $(\xi_{\text{cyclic}}, \overline{\mathcal{P}_\zeta})$-stable.*

Lemma 3.7. (see [NN, Lemma 4.5]) *Let (F, s) be a $(\xi_{\text{cyclic}}, \overline{\mathcal{P}_\zeta})$-stable object, then $F \in \mathcal{P}$.*

Proposition 3.8. (see [NN, Proposition 4.6]) *Let (F, s) be a $(\xi_{\text{cyclic}}, \overline{\mathcal{P}_\zeta})$-stable object, then (F, s) is $(\zeta, \overline{\mathcal{P}})$-stable.*

Remark 3.9. *(1) For a parameter ζ such that $\sum \zeta_l > 0$, we can apply all the argument after a slight modification: first, the information in which chamber $\zeta + d \cdot \mathbf{1}$ $(d \geq 0)$ is contained defines the sequence $k_1, \ldots, k_r$. We define μ_ζ and Ψ_ζ in the same way and denote by $\mathcal{P}_\zeta$ the image of* $(\mu_\zeta(A))$-fmod *under the equivalence $\Psi_\zeta^{-1}[-1]$.*

(2) The Abelian category A-fmod *and the function*

$$Z_{\zeta_{\text{cyclic}}} \colon K(A\text{-fmod}) \simeq \mathbb{Z}^{\hat{I}} \to \mathbb{C}$$

given by

$$Z_{\zeta_{\text{cyclic}}}(\mathbf{v}) := (\mathbf{1} \cdot \mathbf{v}) + (-\zeta_{\text{cyclic}} \cdot \mathbf{v})\sqrt{-1}$$

provide a Bridgeland stability condition $(Z_{\zeta_{\text{cyclic}}}, \{\mathrm{P}(\phi)\}_{\phi \in \mathbb{R}})$, where $\mathrm{P}(\phi)$ is the full subcategory of $D^b(A\text{-fmod})$ of semistable objects with phase ϕ. For $\phi \in \mathbb{R}$, let $\mathcal{P}_{[\phi, \phi+1)}$ denote the full subcategory of objects such that all the factors in their Hardar-Narashimhan filtrations are in

$$\bigcup_{\phi \leq \phi' < \phi+1} \mathrm{P}(\phi').$$

Then we have

$$\mathcal{P}_{[0,1)} = A\text{-fmod}, \quad \mathcal{P}_{[\phi(d), \phi(d)+1)} = \mu_\zeta(A)\text{-fmod}$$

where $\phi(d)$ is determined by the condition $0 \leq \phi(d) < 1/2$ and $\tan(\phi(d)\pi) = d$. This is the reason why our argument works.

(3) Let $A = (Q, \omega)$ be a quiver with a potential which is 3-dimensonal Calabi-Yau and does not have any cycles of length 1 or 2. For a vertex k, we can mutate A at k to get a new one $\mu_k(A)$ and we have a derived equivalence

$$\Psi_k \colon D^b(\mathrm{mod}(A)) \to D^b(\mathrm{mod}(\mu_k(A))).$$

Let $\zeta \in \mathbb{R}^{Q_0}$ be a parameter. Asseme that there exist a sequence $k_1, \dots, k_s$ satisfying the conditions in §3.2. We define the category $\overline{\mathcal{P}_\zeta}$ in the same way. Then the same statement as Proposition 3.6 and 3.8 hold.

3.4 the tilting generator is a vector bundle

In this subsection, we assume that $N \cdot \zeta_k < \sum \zeta_l$ for any $k \neq 0$. Let ν_i $(i \in \tilde{I})$ be real numbers such that

$$\nu_{k+\frac{1}{2}} - \nu_{k-\frac{1}{2}} = N \cdot \zeta_k - \sum \zeta_l$$

for any $k \in \hat{I}$. We denote by $\pi_k \colon \mathbb{R}^{\tilde{I}} \to \mathbb{R}^{\tilde{I}}$ the permutation of the $(k-1/2)$-th and $(k-1/2)$-th elements and set $\pi_s := \pi_{k_s} \circ \cdots \circ \pi_{k_1}$. Note that

$$(\pi_s(\nu))_{k+\frac{1}{2}} - (\pi_s(\nu))_{k-\frac{1}{2}} = N \cdot (\mu_s(\zeta))_k - \sum \zeta_l.$$

By the definition of the sequence (k_s), we have $(\mu_{s-1}(\zeta))_{k_s} > 0$. Recall that we assume $\sum \zeta_l < 0$. Thus we have

$$(\pi_{s-1}(\nu))_{k_s+\frac{1}{2}} < (\pi_{s-1}(\nu))_{k_s-\frac{1}{2}}. \tag{8}$$

Let $\iota_s \colon \tilde{I} \to \tilde{I}$ be the permutation such that the sequence $((\pi_s(\nu))_{\iota_s(i)})_{i \in \tilde{I}}$ is increasing and set $\iota := \iota_r$.

Proposition 3.10. *Under the derived equivalence given in §1, the direct summand $(\mu_\zeta(P))_k$ is a line bundle $\mathcal{L}^\zeta_k$ on Y. Moreover, it is the following map that is associated with the arrow $H^\pm_i$:*

$$\mathcal{L}^\zeta_{i\mp\frac{1}{2}} \hookrightarrow \mathcal{L}^\zeta_{i\mp\frac{1}{2}} \otimes \mathcal{O}_Y(E_{\iota(i)}) \simeq \mathcal{L}^\zeta_{i\pm\frac{1}{2}}.$$

Proof. We prove the claim by induction.

By the argument in the proof of Proposition 1.8, we can verify that the sequence

$$0 \to \mathcal{O} \to \mathcal{O}(E^+_i) \oplus \mathcal{O}(E^-_j) \to \mathcal{O}(E^+_i + E^-_j) \to 0$$

is exact if $i < j$. Thus we have

$$(\mu_s(P))_{k_s} \simeq P_{k_s} \otimes \mathcal{O}(E^+_{\iota_s(k_s-1/2)} + E^-_{\iota_s(k_s+1/2)}).$$

The second half of the claim can be verified by the explicit description of the endomorphism algebra in the proof of Proposition 3.1. □

Corollary 3.11.

$$H^1(Y, (\mathcal{L}^\zeta_k)^{-1}) = 0.$$

Proof. By the argument in the proof of Proposition 3.10, $\mathcal{L}_k^\zeta$ is associated with the divisor which is described as a sum of divisors of the form $E_i^+ + E_j^-$ $(i < j)$. We can verify the claim using the description (5) of H^1 of a line bundle on Y. $\square$

Proposition 3.12. *The set of the torus fixed points $\mathfrak{M}_\zeta(\mathbf{v})^T$ is isolated.*

Proof. By Corollary 3.11 we have $\mathcal{O}_Y \in \mathcal{P}_\zeta$. Let P_ζ be the $\mu_k(A)$-module corresponding to $\mathcal{O}_Y$. By Proposition 3.6 and Proposition 3.8, the moduli space $\mathfrak{M}_\zeta(\mathbf{v})$ parametrizes finite dimensional quotient $\mu_k(A)$-modules V' of P_ζ with $\underline{\dim}\, V' = \mu_k(\mathbf{v})$. Note that

$$(P_\zeta)_k = H^0(Y, (\mathcal{L}_k^\zeta)^{-1})$$

and the T-weight decomposition of H^0 of a line bundle on Y is multiplicity free. Hence the claim follows. $\square$

4 Remarks

In this section, we make some observations on how Kontsevich-Soibelman's wall-crossing formula (9) (they also call the formula (9) by "Factorization Property") would be applied in our setting.

First, we will review the work of Kontsevich-Soibelman ([KS]) very briefly. The core of their work is the construction of the algebra homomorphism from the *"motivic Hall algebra"* to the *"quantum torus"*. For an A_∞-category $\mathcal{C}$, the motivic Hall algebra $H(\mathcal{C})$ is, roughly speaking, the space of motives over the moduli $\mathfrak{Ob}(\mathcal{C})$ of all objects in $\mathcal{C}$, with the product derived from the same diagram as the Ringel-Hall product. The quantum torus is a deformation of a polynomial ring described explicitely in the terms of the numerical datum of $\mathcal{C}$. The homomorphism is given by taking, so to say, weighted Euler characteristics with respect to the *motivic weight*, where the motivic weight is defined using *motivic Milnor fiber* of the potentials coming from the A_∞-structure. The formula (9) is the translation of the Harder-Narashimhan property under this homomorphism.

Remark 4.1. *In the original Donaldson-Thomas invariants defined using symmetric obstruction theory, we adapt the Behrend's function as a weight (see §2.3). It is expected what, after taking the "quasi-classical limit" as $q \to 1$, the motivic weight would coincide with the Behrend's one. In [KS], the proof of the claim for some special situations and some evidences of the claim for more general situations are provided.*

Now, we will explain the statement of "Factorization Property", restricting to our situation. We set $\Lambda := \mathbb{Z}^{\tilde{Q}_0}$ and define the skew symmetric bilinear form $\langle -, - \rangle \colon \Lambda \times \Lambda \to \mathbb{Z}$ by

$$\langle (e_i), (f_i) \rangle := e_\infty \cdot f_0 - e_0 \cdot f_\infty.$$

Let $Z \in \mathrm{Hom}(\Lambda, \mathbb{C})$ be a homomorphism such that $\mathrm{Im}(\mathrm{Z}(\Lambda^+)) > 0$ where $\Lambda^+ = \mathbb{Z}_{\geq 0}^{\tilde{Q}_0}$.

Let D^{μ} be a certain ring of motives including the inverting motives $\mathbb{L}^{-1}$, $[\mathrm{GL}(n)]^{-1}$ $(n \geq 1)$ and the formal symbol $\mathbb{L}^{1/2}$, where $\mathbb{L}$ is the motive of the affine line. We have the homomorphism of rings $\phi\colon D^{\mu} \to \mathbb{Q}(q^{1/2})$ mapping $\mathbb{L}^{1/2}$ to $q^{1/2}$. We define the quantum torus $\mathcal{R}_{\Lambda,q}$ as the $\mathbb{Q}(q^{1/2})$-algebra generated by x_{γ} $(\gamma \in \Lambda)$ with the relation

$$x_{\gamma}x_{\mu} = q^{\frac{1}{2}\langle \gamma,\mu\rangle}x_{\gamma+\mu}.$$

For a strict sector V in the upper half plane, let $\mathcal{C}_V^Z$ denote the category of $\tilde{A}$-modules which can be described as subsequent extensions of Z-semistable objects E such that $Z(E) \in V$. Note that $\mathcal{C}_V^Z$ does not change when Z moves in $\mathrm{Hom}(\Lambda,\mathbb{C})$, unless the values of Z of semistable objects get close to the boundary ∂V. We define an element $A_{V,q}^Z \in \mathcal{R}_{V,q}$ by "weighted" counting of objects in $\mathcal{C}_V^Z$. Informally speaking,

$$A_{V,q}^Z := \sum_{E \in \mathrm{Isom}(\mathcal{C}_V^Z)} \phi\left(\frac{w(E)}{[\mathrm{Aut}(E)]}\right) \cdot x_{\underline{\dim}(E)} \in \mathcal{R}_{V,q},$$

where $w(E) \in D^{\mu}$ is defined by the *motivic Milnor fiber* of the potential of the A_{∞}-algebra algebra $\mathrm{Ext}^*(E,E)$.



Figure 6: V

Assume that V is decomposed into a disjoint union $V = V_1 \sqcup V_2$ in the clockwise order. Then the "Factorization Property" in [KS] claims that

$$A_{V,q}^Z = A_{V_1,q}^Z \cdot A_{V_2,q}^Z. \tag{9}$$

The key fact is, as we mentioned above, the existence of the algebra homomorphism from the motivic Hall algebra to $\mathcal{R}_{V,q}$. Although the category of perverce coherent systems is not Calabi-Yau, Proposition 2.1 would assure the existence of the algebra homomorphism in our case. We can define an element $A_{V,\mathrm{mot}}^Z$ in the motivic Hall algebra and the equation $A_{V,\mathrm{mot}}^Z = A_{V_1,\mathrm{mot}}^Z \cdot A_{V_2,\mathrm{mot}}^Z$ follows from the Harder-Narashimhan property. Now the element $A_{V,q}^Z$ is the image of $A_{V,\mathrm{mot}}^Z$ under the algebra homomorphism.

Now we end up with reviewing and begin to explain how to apply (9) in our setting. Since we are interested in $\tilde{A}$-modules V with $V_{\infty} \simeq \mathbb{C}$, we will work on the quotient algebra

$$\mathcal{R}'_{\Lambda,q} := \mathcal{R}_{\Lambda,q}/(x_{\mathbf{e}})_{\{\mathbf{e}\,|\,e_{\infty}\geq 2\}}.$$

Consider the wall in $\mathrm{Hom}(\Lambda,\mathbb{C})$ such that $\mathbf{e} \in \Lambda^+$ with $e_{\infty} = 0$ and $\mathbf{f} \in \Lambda^+$ with $f_{\infty} = 1$ are send on a same half line in the upper half plane. Assume $\mathbf{e} \in \Lambda$ is primitive (i.e. $\{e_i\}_{i\in Q_0}$ are coprime to each other) and $\mathbf{f} - \mathbf{e} \notin \Lambda^+$.



Figure 7: Λ_0

Let $\Lambda_0 \in \Lambda$ be the sublattice generated by $\mathbf{e}$ and $\mathbf{f}$, l_k be the half line passing through $k \cdot \mathbf{e} + \mathbf{f}$ and l_∞ be the half line passing through $\mathbf{e}$. Take Z^+ and Z^- from the opposite side of the wall so that $l_1, l_2, \ldots, l_\infty$ are mapped on the upper half plane in the clockwise (resp. anticlockwise) way by Z^+ (resp. Z^-). The "Factorization Property" claims

$$A_{l_1}^{Z^+} \cdot A_{l_2}^{Z^+} \cdot \cdots \cdot A_{l_\infty}^{Z^+} = A_{l_\infty}^{Z^-} \cdot \cdots \cdot A_{l_2}^{Z^-} \cdot A_{l_1}^{Z^-}$$

in $\mathcal{R}'_{\Lambda,q}$. We denote

$$\overrightarrow{\prod_k} A_{l_k}^{Z^+} := A_{l_1}^{Z^+} \cdot A_{l_2}^{Z^+} \cdot \cdots, \quad \overleftarrow{\prod_k} A_{l_k}^{Z^-} := \cdots \cdot A_{l_2}^{Z^-} \cdot A_{l_1}^{Z^-}.$$

Note that $A_{l_\infty}^{Z^+} = A_{l_\infty}^{Z^-}$ and we denote this by $A_{l_\infty}^{Z}$. Then the above equation is described as following:

$$\overrightarrow{\prod_k} A_{l_k}^{Z^+} = A_{l_\infty}^{Z} \cdot \left(\overrightarrow{\prod_k} A_{l_k}^{Z^-} \right) \cdot {A_{l_\infty}^{Z}}^{-1}.$$

An element A of $\mathcal{R}'_{\Lambda,q}$ can be uniquely described in the following form:

$$A = \sum_{\mathbf{e};\, e_\infty = 0} (a_{\mathbf{e}}(q) \cdot x_{\mathbf{e}}) + x_\infty \cdot \sum_{\mathbf{e};\, e_\infty = 0} (b_{\mathbf{e}}(q) \cdot x_{\mathbf{e}}).$$

We denote $\sum_{\mathbf{e};\, e_\infty = 0} (b_{\mathbf{e}}(q) \cdot x_{\mathbf{e}})$ by A^{x_∞}. Then

$$(q-1) \cdot \left(\overrightarrow{\prod_k} A_{l_k}^{Z^+} \right)^{x_\infty} \Bigg|_{q=1}$$

makes sense and would coincide with the generating function of virtual counting of the moduli spaces we study in §2. Note that we have

$$\langle k \cdot \mathbf{e} + \mathbf{f}, \mathbf{e} \rangle = e_0.$$

and so

$$x_{m\cdot\mathbf{e}}\cdot x_{k\cdot\mathbf{e}+\mathbf{f}}=q^{m\cdot e_0}x_{k\cdot\mathbf{e}+\mathbf{f}}\cdot x_{m\cdot\mathbf{e}}.$$

Identify $A^{Z+}_{l_\infty}\in\mathcal{R}'_{\Lambda,q}$ with the polynomial in $x_{\mathbf{e}}$, then we have

$$\begin{aligned}\overset{\rightarrow}{\prod_k}A^{Z+}_{l_k}&=A^{Z}_{l_\infty}(x_{\mathbf{e}})\cdot\left(\overset{\leftarrow}{\prod_k}A^{Z-}_{l_k}\right)\cdot A^{Z}_{l_\infty}(x_{\mathbf{e}})^{-1}\\&=\left(\overset{\leftarrow}{\prod_k}A^{Z-}_{l_k}\right)\cdot A^{Z}_{l_\infty}(q^{e_0}x_{\mathbf{e}})\cdot A^{Z}_{l_\infty}(x_{\mathbf{e}})^{-1}\end{aligned}$$

in $\mathcal{R}'_{\Lambda,q}$. Now, if we can compute $A^{Z}_{l_\infty}(q^{e_0}x_{\mathbf{e}})\cdot A^{Z}_{l_\infty}(x_{\mathbf{e}})^{-1}\big|_{q=1}$, we get wall-crossing formulas.

Here again, observations in [KS] will help us. We put $t:=x_{\mathbf{e}}$. Assume that we have the unique simple object E on l_∞ and $\underline{\dim}\,E=\mathbf{e}$. Let B_E be the algebra generated by $\mathrm{Ext}^1(E,E)$ with relations defined from the potential W_E. Then we have

$$A^{Z}_{l_\infty}(qt)=A^{Z}_{l_\infty}(t)\cdot f(t)\tag{10}$$

where $f(t)$ is obtained by counting pairs of cyclic B_E-modules and their cyclic vectors. Applying this formula repeatedly we have

$$A^{Z}_{l_\infty}(q^{e_0}t)\cdot A^{Z}_{l_\infty}(t)^{-1}\big|_{q=1}=\left(f(t)|_{q=1}\right)^{e_0}.$$

Example 4.2. *(1) Assume* $\mathrm{ext}^1(E,E)=0$*. The algebra* B_E *is trivial. Hence we have* $f(t)|_{q=1}=1+t$*. This corresponds to the formula in Theorem 2.14.*

(2) Assume $\mathrm{ext}^1(E,E)=1$ *and* $B_E\simeq\mathbb{C}[z]$*. In this case we have* $f(t)|_{q=1}=(1-t)^{-1}$*. This corresponds to the formula in Theorem 2.18.*

(3) Let us consider the wall corresponding to the imaginary root. The set of simple objects on this wall is $\{\mathcal{O}_y\mid y\in Y\}$*. By the same argument as they show the above equation (10) in [KS], we would have*

$$A^{Z}_{l_\infty}(qt)=A^{Z}_{l_\infty}(t)\cdot f(t)$$

where $f(t)$ *is obtained by counting* 0*-dimensional closed subscheme of* Y*. By the results of [MNOP06] and [BF] we have*

$$f(t)|_{q=1}=M(-t)^{e(Y)}$$

where

$$M(t):=\prod_{n=1}^{\infty}(1-t^n)^{-n}$$

is the MacMahon function. This provides DT-PT correspondence in our situation.

References

[Beh] K. Behrend, *Donaldson-Thomas invariants via microlocal geometry*, math.AG/0507523.

[BF] K. Behrend and B. Fantechi, *Symmetric obstruction theories and Hilbert schemes of points on threefolds*, math.AG/0512556.

[Boc08] R. Bocklandt, *Graded Calabi Yau algebras of dimension* 3, J. Pure Appl. Algebra **212** (2008), no. 1, 14–32.

[Bri02] T. Bridgeland, *Flops and derived categories*, Invent. Math. **147** (2002), no. 3, 613–632.

[CB02] W. Crawley-Boevey, *Decomposition of Marsden-Weinstein reductions for representations of quivers*, Comp. Math. **130** (2002), no. 2, 225–239.

[Ful93] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics studies, Princeton University Press, 1993.

[GV] R. Gopakumar and C. Vafa, *M-theory and topological strings II*, hep-th/9812127.

[HST01] S. Hosono, M. Saito, and A Takahashi, *Relative Lefschetz actions and BPS state counting*, Internat. Math. Res. Notices **15** (2001), 783–816.

[IKP] A. Iqbal and A-K Kashani-Poor, *The vertex on a strip*, hep-th/0410174.

[IU] A. Ishii and K. Ueda, *On moduli spaces of quiver representations associated with brane tilings*, arXiv:0710.1898.

[Joy] D. Joyce, *Configurations in abelian categories IV. Invariants and changing stability conditions*, math/0410268.

[Kin94] A.D. King, *Moduli of renresentations of finite dimensional algebras*, J. Algebra. **45** (1994), no. 4, 515–530.

[KS] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, arXiv:0811.2435.

[Kuz] A. G. Kuznetsov, *Quiver varieties and Hilbert schemes*, AG/0111092.

[LP93] J. Le Potier, *Systèmes cohérents et structures de niveau*, Astérisque (1993), no. 214, 143 pp.

[MNOP06] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory, I*, Comp. Math. **142** (2006), 1263–1285.

[MR] S. Mozgovoy and M. Reineke, *On the noncommutative donaldson-thomas invariants arising from brane tilings*, arXiv:0809.0117.

[Nak94] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. **76** (1994), no. 2, 365–416.

[Nak98] ______, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. 3, 515–560.

[Nak99] ______, *Lectures on Hilbert schemes of points on surfaces*, University Lecture Series, American Mathematical Society, Providence, RI, 1999.

[Nak01] ______, *Quiver varieties and finite-dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), no. 1, 145–238.

[NN] K. Nagao and H. Nakajima, *Counting invarinats of perverse coherent systems on 3-folds and their wall-crossings*, arXiv:0809.2992.

[NYa] H. Nakajima and K. Yoshioka, *Perverse coherent sheaves on blow-up. I. a quiver description*, arXiv:0802.3120.

[NYb] ______, *Perverse coherent sheaves on blow-up. II. wall-crossing and Betti numbers formula*, arXiv:0806.0463.

[PTa] R. Pandharipande and R.P. Thomas, *Curve counting via stable pairs in the derived category*, arXiv:0707.2348.

[PTb] ______, *Stable pairs and BPS invariants*, arXiv:0711.3899.

[Rud97] A. Rudakov, *Stability for an Abelian category*, J. Algebra. **197** (1997), no. 1, 231–245.

[Sze] B. Szendroi, *Non-commutative Donaldson-Thomas theory and the conifold*, arXiv:0705.3419v3.

[Tho00] R. P. Thomas, *A holomorphic Casson invariant for Calabi-Yau* 3*-folds, and bundles on* $K3$ *fibrations*, J. Differential Geom. **54** (2000), no. 2, 367–438.

[Tod] Y. Toda, *Birational Calabi-Yau* 3*-folds and BPS state counting*, arXiv:0707.1643.

[VdB] M. Van den Bergh, *Non-commutative crepant resolutions*, math/0211064.

[VdB04] ______, *Three-dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no. 3, 423–455.

[Youa] B. Young, *Computing a pyramid partition generating function with dimer shuffling*, arXiv:0709.3079.

[Youb] J Young, B. with an appendix by Bryan, *Generating functions for colored 3d Young diagrams and the Donaldson-Thomas invariants of orbifolds*, arXiv:0802.3948.

Kentaro Nagao
Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan
kentaron@kurims.kyoto-u.ac.jp